

**FINAL JEE-MAIN EXAMINATION – APRIL, 2024**

 (Held On Thursday 04<sup>th</sup> April, 2024)

TIME : 9 : 00 AM to 12 : 00 NOON

**CHEMISTRY**
**TEST PAPER WITH ANSWER**
**SECTION-A**





61. What pressure (bar) of  $H_2$  would be required to make emf of hydrogen electrode zero in pure water at  $25^\circ C$  ?  
 (1)  $10^{-14}$  (2)  $10^{-7}$  (3) 1 (4) 0.5

**Allen Ans. (1)**
**NTA Ans. (3)**

62. The correct sequence of ligands in the order of decreasing field strength is :  
 (1)  $CO > H_2O > F^- > S^{2-}$   
 (2)  $^-OH > F^- > NH_3 > CN^-$   
 (3)  $NCS^- > EDTA^{4-} > CN^- > CO$   
 (4)  $S^{2-} > ^-OH > EDTA^{4-} > CO$

**Ans. (1)**

63. Match List -I with List II:

	List - I Mechanism steps	List - II Effect
(A)		(I) - E effect
(B)		(II) - R effect
(C)		(III) + E effect
(D)		(IV) + R effect

Choose the **correct** answer from the options given below :

- (1) (A) – (IV), (B) – (III), (C) – (I), (D) – (II)  
 (2) (A) – (III), (B) – (I), (C) – (II), (D) – (IV)  
 (3) (A) – (II), (B) – (IV), (C) – (III), (D) – (I)  
 (4) (A) – (I), (B) – (II), (C) – (IV), (D) – (III)

**Ans. (1)**

64. What will be the decreasing order of basic strength of the following conjugate bases ?




- (1)  $C\bar{I} > OH > R\bar{O} > CH_3CO\bar{O}$   
 (2)  $R\bar{O} > ^-OH > CH_3CO\bar{O} > C\bar{I}$   
 (3)  $^-OH > R\bar{O} > CH_3CO\bar{O} > C\bar{I}$   
 (4)  $C\bar{I} > R\bar{O} > ^-OH > CH_3CO\bar{O}$









**Ans. (2)**

65. In the precipitation of the iron group (III) in qualitative analysis, ammonium chloride is added before adding ammonium hydroxide to :

- (1) prevent interference by phosphate ions  
 (2) decrease concentration of  $^-OH$  ions  
 (3) increase concentration of  $Cl^-$  ions  
 (4) increase concentration of  $NH_4^+$  ions

**Ans. (2)**


Identify (B) and (C) and how are (A) and (C) related ?

	(B)	(C)	
(1)			functional group isomers
(2)			Derivative
(3)			position isomers
(4)			chain isomers

**Ans. (3)**


Download the new **ALLEN app**  
 & enroll for **Online Programs**


CLICK HERE TO  
 DOWNLOAD

67. One of the commonly used electrode is calomel electrode. Under which of the following categories calomel electrode comes ?

- (1) Metal – Insoluble Salt – Anion electrodes
- (2) Oxidation – Reduction electrodes
- (3) Gas – Ion electrodes
- (4) Metal ion – Metal electrodes

Ans. (1)

68. Number of complexes from the following with even number of unpaired "d" electrons is \_\_\_\_.



[Given atomic numbers : V = 23, Cr = 24, Fe = 26, Ni = 28, Cu = 29]

- (1) 2
- (2) 4
- (3) 5
- (4) 1


Ans. (1)

69. Which one of the following molecules has maximum dipole moment ?

- (1)  $\text{NF}_3$
- (2)  $\text{CH}_4$
- (3)  $\text{NH}_3$
- (4)  $\text{PF}_5$

Ans. (3)

70. Number of molecules/ions from the following in which the central atom is involved in  $\text{sp}^3$  hybridization is \_\_\_\_\_.



- (1) 2
- (2) 4
- (3) 3
- (4) 1

Ans. (1)

71. Which among the following is incorrect statement?

- (1) Electromeric effect dominates over inductive effect
- (2) The electromeric effect is, temporary effect
- (3) The organic compound shows electromeric effect in the presence of the reagent only
- (4) Hydrogen ion ( $\text{H}^+$ ) shows negative electromeric effect

Ans. (4)

72. Given below are two statements :

**Statement I :** Acidity of  $\alpha$ -hydrogens of aldehydes and ketones is responsible for Aldol reaction.

**Statement II :** Reaction between benzaldehyde and ethanal will NOT give Cross – Aldol product.

In the light of above statements, choose the **most appropriate** answer from the options given below.

- (1) Both **Statement I** and **Statement II** are correct.
- (2) Both **Statement I** and **Statement II** are incorrect.
- (3) **Statement I** is incorrect but **Statement II** is correct.
- (4) **Statement I** is correct but **Statement II** is incorrect.



Ans. (4)

73. Which of the following nitrogen containing compound does not give Lassaigne's test ?

- (1) Phenyl hydrazine
- (2) Glycine
- (3) Urea
- (4) Hydrazine

Ans. (4)

74. Which of the following is the correct structure of L-Glucose ?




Ans. (1)

75. The element which shows only one oxidation state other than its elemental form is :

- (1) Cobalt
- (2) Scandium
- (3) Titanium
- (4) Nickel


Ans. (2)



Download the new ALLEN app  
& enroll for Online Programs

CLICK HERE TO  
DOWNLOAD

76. Identify the product in the following reaction :



Ans. (4)

77. Number of elements from the following that CANNOT form compounds with valencies which match with their respective group valencies is \_\_\_\_\_.

B, C, N, S, O, F, P, Al, Si  
 (1) 7 (2) 5 (3) 6 (4) 3


Ans. (4)

78. The Molarity (M) of an aqueous solution containing 5.85 g of NaCl in 500 mL water is :  
 (Given : Molar Mass Na : 23 and Cl : 35.5  $\text{gmol}^{-1}$ )

(1) 20 (2) 0.2  
 (3) 2 (4) 4

Ans. (2)

79. Identify the correct set of reagents or reaction conditions 'X' and 'Y' in the following set of transformation.



(1) X = conc.alc. NaOH,  $80^\circ\text{C}$ , Y =  $\text{Br}_2/\text{CHCl}_3$   
 (2) X = dil.aq. NaOH,  $20^\circ\text{C}$ , Y = HBr/acetic acid  
 (3) X = conc.alc. NaOH,  $80^\circ\text{C}$ , Y = HBr/acetic acid  
 (4) X = dil.aq. NaOH,  $20^\circ\text{C}$ , Y =  $\text{Br}_2/\text{CHCl}_3$

Ans. (3)

80. The correct order of first ionization enthalpy values of the following elements is :

(A) O (B) N  
 (C) Be (D) F  
 (E) B

Choose the correct answer from the options given below :

(1)  $\text{B} < \text{D} < \text{C} < \text{E} < \text{A}$  (2)  $\text{E} < \text{C} < \text{A} < \text{B} < \text{D}$   
 (3)  $\text{C} < \text{E} < \text{A} < \text{B} < \text{D}$  (4)  $\text{A} < \text{B} < \text{D} < \text{C} < \text{E}$


Ans. (2)

SECTION-B

81. The enthalpy of formation of ethane ( $\text{C}_2\text{H}_6$ ) from ethylene by addition of hydrogen where the bond-energies of C – H, C – C, H – H are 414 kJ, 347 kJ, 615 kJ and 435 kJ respectively is - \_\_\_\_\_ kJ.

Ans. (125)

82. The number of correct reaction(s) among the following is \_\_\_\_\_.



Ans. (1)

83. X g of ethylamine is subjected to reaction with  $\text{NaNO}_2/\text{HCl}$  followed by water; evolved dinitrogen gas which occupied 2.24 L volume at STP.

X is \_\_\_\_\_  $\times 10^{-1}$  g.

Ans. (45)

84. The de-Broglie's wavelength of an electron in the 4<sup>th</sup> orbit is \_\_\_\_\_  $\pi a_0$ . ( $a_0$  = Bohr's radius)

Ans. (8)

85. Only 2 mL of  $\text{KMnO}_4$  solution of unknown molarity is required to reach the end point of a titration of 20 mL of oxalic acid (2 M) in acidic medium. The molarity of  $\text{KMnO}_4$  solution should be \_\_\_\_\_ M.

Allen Ans. (8)


NTA Ans. (50)


 Download the new ALLEN app & enroll for Online Programs
 [CLICK HERE TO DOWNLOAD](#)

86. Consider the following reaction  
 $\text{MnO}_2 + \text{KOH} + \text{O}_2 \rightarrow \text{A} + \text{H}_2\text{O}$ .  
 Product 'A' in neutral or acidic medium disproportionate to give products 'B' and 'C' along with water. The sum of spin-only magnetic moment values of B and C is \_\_\_\_\_ BM. (nearest integer)  
 (Given atomic number of Mn is 25)

Ans. (4)

87. Consider the following transformation involving first order elementary reaction in each step at constant temperature as shown below.



Some details of the above reaction are listed below.

Step	Rate constant (sec <sup>-1</sup> )	Activation energy (kJ mol <sup>-1</sup> )
1	$k_1$	300
2	$k_2$	200
3	$k_3$	$E_{a_3}$

If the overall rate constant of the above transformation (k) is given as  $k = \frac{k_1 k_2}{k_3}$  and the

overall activation energy ( $E_a$ ) is 400 kJ mol<sup>-1</sup>, then the value of  $E_{a_3}$  is \_\_\_\_\_ kJ mol<sup>-1</sup> (nearest integer)

Ans. (100)

88. 2.5 g of a non-volatile, non-electrolyte is dissolved in 100 g of water at 25°C. The solution showed a boiling point elevation by 2°C. Assuming the solute concentration is negligible with respect to the solvent concentration, the vapour pressure of the resulting aqueous solution is \_\_\_\_\_ mm of Hg (nearest integer)

[Given : Molal boiling point elevation constant of water ( $K_b$ ) = 0.52 K. kg mol<sup>-1</sup>,

1 atm pressure = 760 mm of Hg, molar mass of water = 18 g mol<sup>-1</sup>]

Ans. (707)

89. The number of different chain isomers for C<sub>7</sub>H<sub>16</sub> is \_\_\_\_\_.

Ans. (9)

90. Number of molecules/species from the following having one unpaired electron is \_\_\_\_\_.

O<sub>2</sub>, O<sub>2</sub><sup>-1</sup>, NO, CN<sup>-1</sup>, O<sub>2</sub><sup>2-</sup>

Ans. (2)



Download the new ALLEN app  
 & enroll for Online Programs

CLICK HERE TO  
 DOWNLOAD



Are you targeting **JEE 2025** ?

Ace it with **ALLEN's**  
**Leader Course**

Online Program 18 APRIL '24

Offline Program 24 APRIL '24



**ALLEN**


Get The Latest  
**IIT-JEE** Special Books  
at Your Door Steps...!!

**JOIN THE JOURNEY OF LEARNING**

*with*

**SCORE TEST PAPERS | HANDBOOKS**  
**JEE-MAIN PYQ's | JEE-Adv. PYQ's**

**SHOP NOW**



Available in  
**HINDI & ENGLISH**