

**FINAL JEE-MAIN EXAMINATION – APRIL, 2024**

**(Held On Tuesday 09<sup>th</sup> April, 2024)**

**TIME : 3 : 00 PM to 6 : 00 PM**

**PHYSICS**

**TEST PAPER WITH ANSWER**


**SECTION-A**

31. A nucleus at rest disintegrates into two smaller nuclei with their masses in the ratio of 2:1. After disintegration they will move :-

- (1) In opposite directions with speed in the ratio of 1:2 respectively
- (2) In opposite directions with speed in the ratio of 2:1 respectively
- (3) In the same direction with same speed.
- (4) In opposite directions with the same speed.

Ans. (1)

32. The following figure represents two biconvex lenses  $L_1$  and  $L_2$  having focal length 10 cm and 15 cm respectively. The distance between  $L_1$  &  $L_2$  is :



- (1) 10 cm
- (2) 15 cm
- (3) 25 cm
- (4) 35 cm

Ans. (3)

33. The temperature of a gas is  $-78^\circ\text{C}$  and the average translational kinetic energy of its molecules is  $K$ . The temperature at which the average translational kinetic energy of the molecules of the same gas becomes  $2K$  is :

- (1)  $-39^\circ\text{C}$
- (2)  $117^\circ\text{C}$
- (3)  $127^\circ\text{C}$
- (4)  $-78^\circ\text{C}$

Ans. (2)

34. A hydrogen atom in ground state is given an energy of 10.2 eV. How many spectral lines will be emitted due to transition of electrons ?

- (1) 6
- (2) 3
- (3) 10
- (4) 1


Ans. (4)

35. The magnetic field in a plane electromagnetic wave is  $B_y = (3.5 \times 10^{-7}) \sin(1.5 \times 10^3 x + 0.5 \times 10^{11} t)$  T. The corresponding electric field will be

- (1)  $E_y = 1.17 \sin(1.5 \times 10^3 x + 0.5 \times 10^{11} t) \text{Vm}^{-1}$
- (2)  $E_z = 105 \sin(1.5 \times 10^3 x + 0.5 \times 10^{11} t) \text{Vm}^{-1}$
- (3)  $E_z = 1.17 \sin(1.5 \times 10^3 x + 0.5 \times 10^{11} t) \text{Vm}^{-1}$
- (4)  $E_y = 10.5 \sin(1.5 \times 10^3 x + 0.5 \times 10^{11} t) \text{Vm}^{-1}$

Ans. (2)

36. A square loop of side 15 cm being moved towards right at a constant speed of 2 cm/s as shown in figure. The front edge enters the 50 cm wide magnetic field at  $t = 0$ . The value of induced emf in the loop at  $t = 10$  s will be :




- (1) 0.3 mV
- (2) 4.5 mV
- (3) zero
- (4) 3 mV

Ans. (3)

37. Two cars are travelling towards each other at speed of  $20 \text{ m s}^{-1}$  each. When the cars are 300 m apart, both the drivers apply brakes and the cars retard at the rate of  $2 \text{ m s}^{-2}$ . The distance between them when they come to rest is :

- (1) 200 m
- (2) 50 m
- (3) 100 m
- (4) 25 m


Ans. (3)



Download the new **ALLEN app** & enroll for **Online Programs**

CLICK HERE TO DOWNLOAD

38. The  $I$ - $V$  characteristics of an electronic device shown in the figure. The device is :



- (1) a solar cell
- (2) a transistor which can be used as an amplifier
- (3) a zener diode which can be used as voltage regulator
- (4) a diode which can be used as a rectifier

Ans. (3)

39. The excess pressure inside a soap bubble is thrice the excess pressure inside a second soap bubble. The ratio between the volume of the first and the second bubble is :

- (1) 1 : 9
- (2) 1 : 3
- (3) 1 : 81
- (4) 1 : 27

Ans. (4)

40. The de-Broglie wavelength associated with a particle of mass  $m$  and energy  $E$  is  $h / \sqrt{2mE}$ . The dimensional formula for Planck's constant is :

- (1)  $[ML^{-1}T^{-2}]$
- (2)  $[ML^2T^{-1}]$
- (3)  $[MLT^{-2}]$
- (4)  $[M^2L^2T^{-2}]$


Ans. (2)

41. A satellite of  $10^3$  kg mass is revolving in circular orbit of radius  $2R$ . If  $\frac{10^4 R}{6}$  J energy is supplied to the satellite, it would revolve in a new circular orbit of radius :

- (use  $g = 10\text{m/s}^2$ ,  $R = \text{radius of earth}$ )
- (1)  $2.5 R$
  - (2)  $3 R$
  - (3)  $4 R$
  - (4)  $6 R$

Ans. (4)


42. The effective resistance between  $A$  and  $B$ , if resistance of each resistor is  $R$ , will be



- (1)  $\frac{2}{3} R$
- (2)  $\frac{8R}{3}$
- (3)  $\frac{5R}{3}$
- (4)  $\frac{4R}{3}$

Ans. (2)

43. Five charges  $+q$ ,  $+5q$ ,  $-2q$ ,  $+3q$  and  $-4q$  are situated as shown in the figure. The electric flux due to this configuration through the surface  $S$  is :



- (1)  $\frac{5q}{\epsilon_0}$
- (2)  $\frac{4q}{\epsilon_0}$
- (3)  $\frac{3q}{\epsilon_0}$
- (4)  $\frac{q}{\epsilon_0}$

Ans. (2)

44. A proton and a deuteron ( $q = +e$ ,  $m = 2.0u$ ) having same kinetic energies enter a region of uniform magnetic field  $\vec{B}$ , moving perpendicular to  $\vec{B}$ . The ratio of the radius  $r_d$  of deuteron path to the radius  $r_p$  of the proton path is :

- (1) 1 : 1
- (2) 1 :  $\sqrt{2}$
- (3)  $\sqrt{2} : 1$
- (4) 1 : 2

Ans. (3)

45. UV light of 4.13 eV is incident on a photosensitive metal surface having work function 3.13 eV. The maximum kinetic energy of ejected photoelectrons will be :

- (1) 4.13 eV                      (2) 1 eV  
 (3) 3.13 eV                      (4) 7.26 eV


Ans. (2)

46. The energy released in the fusion of 2 kg of hydrogen deep in the sun is  $E_H$  and the energy released in the fission of 2 kg of  $^{235}\text{U}$  is  $E_U$ . The ratio  $\frac{E_H}{E_U}$  is approximately :

- (Consider the fusion reaction as  $4\text{}^1_1\text{H} + 2\text{e}^- \rightarrow \text{}^4_2\text{He} + 2\nu + 6\gamma + 26.7\text{MeV}$ , energy released in the fission reaction of  $^{235}\text{U}$  is 200 MeV per fission nucleus and  $N_A = 6.023 \times 10^{23}$ )
- (1) 9.13                              (2) 15.04  
 (3) 7.62                              (4) 25.6

Ans. (3)


47. A real gas within a closed chamber at 27°C undergoes the cyclic process as shown in figure. The gas obeys  $PV^3 = RT$  equation for the path A to B. The net work done in the complete cycle is (assuming  $R = 8\text{J/molK}$ ):



- (1) 225 J                              (2) 205 J  
 (3) 20 J                                (4) -20 J

Ans. (2)

48. A 1 kg mass is suspended from the ceiling by a rope of length 4m. A horizontal force 'F' is applied at the mid point of the rope so that the rope makes an angle of 45° with respect to the vertical axis as shown in figure. The magnitude of F is :




- (1)  $\frac{10}{\sqrt{2}}$  N                      (2) 1 N  
 (3)  $\frac{1}{10 \times \sqrt{2}}$  N                      (4) 10 N

Ans. (4)

49. A spherical ball of radius  $1 \times 10^{-4}$  m and density  $10^5 \text{ kg/m}^3$  falls freely under gravity through a distance  $h$  before entering a tank of water. If after entering in water the velocity of the ball does not change, then the value of  $h$  is approximately :

- (The coefficient of viscosity of water is  $9.8 \times 10^{-6} \text{ N s/m}^2$ )
- (1) 2296 m                              (2) 2249 m  
 (3) 2518 m                              (4) 2396 m

Ans. (3)



In the truth table of the above circuit the value of X and Y are :

- (1) 1, 1                                      (2) 1, 0  
 (3) 0, 1                                      (4) 0, 0

Ans. (1)

SECTION-B

51. A straight magnetic strip has a magnetic moment of  $44 \text{ Am}^2$ . If the strip is bent in a semicircular shape, its magnetic moment will be .....  $\text{Am}^2$

(Given  $\pi = \frac{22}{7}$ )

Ans. (28)

52. A particle of mass  $0.50 \text{ kg}$  executes simple harmonic motion under force  $F = -50(\text{Nm}^{-1})x$ . The time period of oscillation is  $\frac{x}{35} \text{ s}$ . The value of  $x$  is .....


(Given  $\pi = \frac{22}{7}$ )

Ans. (22)

53. A capacitor of reactance  $4\sqrt{3}\Omega$  and a resistor of resistance  $4\Omega$  are connected in series with an ac source of peak value  $8\sqrt{2} \text{ V}$ . The power dissipation in the circuit is .....  $\text{W}$ .

Ans. (4)

54. An electric field  $\vec{E} = (2x\hat{i})\text{NC}^{-1}$  exists in space. A cube of side  $2\text{m}$  is placed in the space as per figure given below. The electric flux through the cube is .....  $\text{Nm}^2/\text{C}$ .




Ans. (16)

55. A circular disc reaches from top to bottom of an inclined plane of length  $l$ . When it slips down the plane, it takes  $t$  s. When it rolls down the plane then it takes  $\left(\frac{\alpha}{2}\right)^{1/2} t$  s, where  $\alpha$  is .....

Ans. (3)

56. To determine the resistance ( $R$ ) of a wire, a circuit is designed below, The V-I characteristic curve for this circuit is plotted for the voltmeter and the ammeter readings as shown in figure. The value of  $R$  is .....  $\Omega$ .



Ans. (2500)

57. The resultant of two vectors  $\vec{A}$  and  $\vec{B}$  is perpendicular to  $\vec{A}$  and its magnitude is half that of  $\vec{B}$ . The angle between vectors  $\vec{A}$  and  $\vec{B}$  is .....

Ans. (150)

58. Monochromatic light of wavelength  $500 \text{ nm}$  is used in Young's double slit experiment. An interference pattern is obtained on a screen. When one of the slits is covered with a very thin glass plate (refractive index =  $1.5$ ), the central maximum is shifted to a position previously occupied by the 4<sup>th</sup> bright fringe. The thickness of the glass-plate is .....  $\mu\text{m}$ .


Ans. (4)

59. A force  $(3x^2 + 2x - 5) \text{ N}$  displaces a body from  $x = 2 \text{ m}$  to  $x = 4\text{m}$ . Work done by this force is .....  $\text{J}$ .

Ans. (58)

60. At room temperature ( $27^\circ\text{C}$ ), the resistance of a heating element is  $50\Omega$ . The temperature coefficient of the material is  $2.4 \times 10^{-4} \text{ }^\circ\text{C}^{-1}$ . The temperature of the element, when its resistance is  $62 \Omega$ , is .....  $^\circ\text{C}$ .

Ans. (1027)



Download the new ALLEN app & enroll for Online Programs

CLICK HERE TO DOWNLOAD

Are you targeting **JEE 2025** ?

Ace it with **ALLEN's**  
**Leader Course**

Online Program 18 APRIL '24

Offline Program 24 APRIL '24



**ALLEN**


Get The Latest  
**IIT-JEE Special Books**  
at Your Door Steps...!!

**JOIN THE JOURNEY OF LEARNING**

with

**SCORE TEST PAPERS | HANDBOOKS**  
**JEE-MAIN PYQ's | JEE-Adv. PYQ's**

**SHOP NOW**



Available in  
**HINDI & ENGLISH**