

**FINAL JEE-MAIN EXAMINATION – APRIL, 2024**

(Held On Tuesday 09<sup>th</sup> April, 2024)


TIME : 9 : 00 AM to 12 : 00 NOON

**CHEMISTRY**

**TEST PAPER WITH SOLUTION**

**SECTION-A**

61. The molar conductivity for electrolytes A and B are plotted against  $C^{1/2}$  as shown below. Electrolytes A and B respectively are :



**A**

**B**

- |                        |                    |
|------------------------|--------------------|
| (1) Weak electrolyte   | weak electrolyte   |
| (2) Strong electrolyte | strong electrolyte |
| (3) Weak electrolyte   | strong electrolyte |
| (4) Strong electrolyte | weak electrolyte   |

Ans. (3)


Sol. A → Weak electrolyte  
B → Strong electrolyte

62. Methods used for purification of organic compounds are based on :
- (1) neither on nature of compound nor on the impurity present.
  - (2) nature of compound only.
  - (3) nature of compound and presence of impurity.
  - (4) presence of impurity only.

Ans. (3)

Sol. Organic compounds are purified based on their nature and impurity present in it.


63. In the following sequence of reaction, the major products B and C respectively are :



- (1) D-C<sub>6</sub>H<sub>4</sub>-D and F-C<sub>6</sub>H<sub>4</sub>-F
- (2) D-C<sub>6</sub>H<sub>4</sub>-D and F-C<sub>6</sub>H<sub>4</sub>-F
- (3) D-C<sub>6</sub>H<sub>4</sub>-D and F-C<sub>6</sub>H<sub>4</sub>-F
- (4) D-C<sub>6</sub>H<sub>4</sub>-D and F-C<sub>6</sub>H<sub>4</sub>-F

Ans. (1)

Sol.



64. Correct order of basic strength of Pyrrole ,

Pyridine and Piperidine is:

- (1) Piperidine > Pyridine > Pyrrole
- (2) Pyrrole > Pyridine > Piperidine
- (3) Pyridine > Piperidine > Pyrrole
- (4) Pyrrole > Piperidine > Pyridine

Ans. (1)

Sol. Order of basic strength is  
N(sp<sup>3</sup>, localized lone pair) > N(sp<sup>2</sup>, localized lone pair) > N(sp<sup>2</sup>, delocalized lone pair, aromatic)  
∴ Piperidine > Pyridine > Pyrrole

65. In which one of the following pairs the central atoms exhibit sp<sup>2</sup> hybridization ?

- (1) BF<sub>3</sub> and NO<sub>2</sub><sup>-</sup>
- (2) NH<sub>2</sub><sup>-</sup> and H<sub>2</sub>O
- (3) H<sub>2</sub>O and NO<sub>2</sub>
- (4) NH<sub>2</sub><sup>-</sup> and BF<sub>3</sub>

Ans. (1)

Sol. BF<sub>3</sub> → sp<sup>2</sup>  
NO<sub>2</sub><sup>-</sup> → sp<sup>2</sup>  
H<sub>2</sub>O → sp<sup>3</sup>  
NO<sub>2</sub> → sp<sup>2</sup>  
NH<sub>2</sub><sup>-</sup> → sp<sup>3</sup>



Download the new ALLEN app  
& enroll for Online Programs

CLICK HERE TO  
DOWNLOAD


66. The  $F^-$  ions make the enamel on teeth much harder by converting hydroxyapatite (the enamel on the surface of teeth) into much harder fluoroapatite having the formula.

- (1)  $[3(Ca_3(PO_4)_2).CaF_2]$
- (2)  $[3(Ca_2(PO_4)_2).Ca(OH)_2]$
- (3)  $[3(Ca_3(PO_4)_3).CaF_2]$
- (4)  $[3(Ca_3(PO_4)_2).Ca(OH)_2]$

Ans. (1)

Sol. Fluoroapatite  $\Rightarrow [3Ca_3(PO_4)_2.CaF_2]$

67. Relative stability of the contributing structures is :



- (1) (I) > (III) > (II)
- (2) (I) > (II) > (III)
- (3) (II) > (I) > (III)
- (4) (III) > (II) > (I)

Ans. (2)

Sol. (1) Neutral structures are more stable than charged ones. Therefore I is more stable than II and III.

(2) +ve charge on less electronegative atom is more stable i.e.,  $C^\oplus$  is more stable than  $O^\oplus$

$\therefore$  Order is I > II > III

68. Given below are two statements :

**Statement (I)** : The oxidation state of an element in a particular compound is the charge acquired by its atom on the basis of electron gain enthalpy consideration from other atoms in the molecule.

**Statement (II)** :  $p\pi-p\pi$  bond formation is more prevalent in second period elements over other periods.

In the light of the above statements, choose the **most appropriate** answer from the options given below :

- (1) Both **Statement I** and **Statement II** are incorrect
- (2) **Statement I** is correct but **Statement II** is incorrect
- (3) Both **Statement I** and **Statement II** are correct
- (4) **Statement I** is incorrect but **Statement II** is correct

Ans. (4)

Sol. Oxidation state of an element in a particular compound is defined by the charge acquired by its atom on the basis of electronegativity consideration from other atoms in molecule.

69. Given below are two statements : one is labelled as **Assertion (A)** and the other is labelled as **Reason (R)** :

**Assertion (A)** :  $S_N2$  reaction of  $C_6H_5CH_2Br$  occurs more readily than the  $S_N2$  reaction of  $CH_3CH_2Br$ .


**Reason (R)** : The partially bonded unhybridized p-orbital that develops in the trigonal bipyramidal transition state is stabilized by conjugation with the phenyl ring.

In the light of the above statements, choose the **most appropriate** answer from the options given below :






- (1) **(A)** is not correct but **(R)** is correct
- (2) Both **(A)** and **(R)** are correct but **(R)** is not the correct explanation of **(A)**
- (3) Both **(A)** and **(R)** are correct and **(R)** is the correct explanation of **(A)**
- (4) **(A)** is correct but **(R)** is not correct

Ans. (3)

Sol. The benzyl group acts in much the same way using the  $\pi$ -system of the benzene ring for conjugation with the p-orbital in the transition state.



70. For the given compounds, the correct order of increasing  $pK_a$  value :




- (1) (E) < (D) < (C) < (B) < (A)  
 (2) (D) < (E) < (C) < (B) < (A)  
 (3) (E) < (D) < (B) < (A) < (C)  
 (4) (B) < (D) < (A) < (C) < (E)


Ans. **BONUS**

NTA Ans. (4)

Sol. Acidic strength order :-



Correct  $pK_a$  Order :



All options are incorrect.

71. Given below are two statements : one is labelled as **Assertion (A)** : and the other is labelled as **Reason (R)**.

**Assertion (A)** : Both rhombic and monoclinic sulphur exist as  $S_8$  while oxygen exists as  $O_2$ .

**Reason (R)** : Oxygen forms  $p\pi-p\pi$  multiple bonds with itself and other elements having small size and high electronegativity like C, N, which is not possible for sulphur.


In the light of the above statements, choose the **most appropriate** answer from the options given below :

- (1) Both (A) and (R) are correct and (R) is the correct explanation of (A).  
 (2) Both (A) and (R) are correct but (R) is not the correct explanation of (A).  
 (3) (A) is correct but (R) is not correct.  
 (4) (A) is not correct but (R) is correct.

Ans. (3)

Sol. Oxygen can form  $2p\pi-2p\pi$  multiple bond with itself due to its small size while sulphur cannot form multiple bond with itself as  $3p\pi-3p\pi$  bond will be unstable due to large size of sulphur, but sulphur can form multiple bond with small size atom like C and N.

eg.  $S=C=S$



72. Given below are two statements : one is labelled as **Assertion (A)** and the other is labelled as **Reason (R)**.

**Assertion (A)**: The total number of geometrical isomers shown by  $[Co(en)_2Cl_2]^+$  complex ion is three


**Reason (R)**:  $[Co(en)_2Cl_2]^+$  complex ion has an octahedral geometry.

In the light of the above statements, choose the **most appropriate** answer from the options given below :

- (1) Both (A) and (R) are correct and (R) is the correct explanation of (A).  
 (2) (A) is correct but (R) is not correct.  
 (3) (A) is not correct but (R) is correct.  
 (4) Both (A) and (R) are correct but (R) is not the correct explanation of (A).

Ans. (3)

Sol.  $[Co(en)_2Cl_2]^+$  has octahedral geometry with two geometrical isomers.




73. The electronic configuration of Cu(II) is  $3d^9$  whereas that of Cu(I) is  $3d^{10}$ . Which of the following is correct ?

- (1) Cu(II) is less stable  
 (2) Stability of Cu(I) and Cu(II) depends on nature of copper salts  
 (3) Cu(II) is more stable  
 (4) Cu(I) and Cu(II) are equally stable


Ans. (3)

Sol. Cu(II) is more stable than Cu(I) because hydration energy of  $Cu^{+2}$  ion compensate  $IE_2$  of Cu.







Download the new **ALLEN** app  
 & enroll for **Online Programs**



CLICK HERE TO  
 DOWNLOAD



What is the structure of C ?

- (1) 
- (2) 
- (3) 
- (4) 

Ans. (1)



75. Compare the energies of following sets of quantum numbers for multielectron system.

- (A)  $n = 4, l = 1$       (B)  $n = 4, l = 2$   
 (C)  $n = 3, l = 1$       (D)  $n = 3, l = 2$   
 (E)  $n = 4, l = 0$

Choose the correct answer from the options given below :

- (1) (B) > (A) > (C) > (E) > (D)  
 (2) (E) > (C) < (D) < (A) < (B)  
 (3) (E) > (C) > (A) > (D) > (B)  
 (4) (C) < (E) < (D) < (A) < (B)

Ans. (4)


Sol. Energy level can be determined by comparing  $(n + l)$  values





- (A)  $n = 4, l = 1 \Rightarrow (n + l) = 5$   
 (B)  $n = 4, l = 2 \Rightarrow (n + l) = 6$   
 (C)  $n = 3, l = 1 \Rightarrow (n + l) = 4$   
 (D)  $n = 3, l = 2 \Rightarrow (n + l) = 5$   
 (E)  $n = 4, l = 0 \Rightarrow (n + l) = 4$

For same value of  $(n + l)$ , orbital having higher value of  $n$ , will have more energy.

(B) > (A) > (D) > (E) > (C)


76. Identify major product "X" formed in the following reaction :




- (1) 
- (2) 
- (3) 
- (4) 

Ans. (3)

Sol. This is Gattermann-Koch reaction




77. Identify the product A and product B in the following set of reactions.



- (1) A- $\text{CH}_3\text{CH}_2\text{CH}_2\text{-OH}$ , B- $\text{CH}_3\text{CH}_2\text{CH}_2\text{-OH}$   
 (2) A- $\text{CH}_3\text{CH}_2\text{CH}_2\text{-OH}$ , B- $\text{CH}_3\text{CH}(\text{OH})\text{-CH}_3$   
 (3) A- $\text{CH}_3\text{-CH}(\text{OH})\text{-CH}_3$ , B- $\text{CH}_3\text{CH}_2\text{CH}_2\text{-OH}$   
 (4) A- $\text{CH}_3\text{CH}_2\text{CH}_3$ , B- $\text{CH}_3\text{CH}_2\text{CH}_3$



Ans. (3)






Download the new ALLEN app  
& enroll for Online Programs

CLICK HERE TO  
DOWNLOAD

Sol. (1) Hydration Reaction :




(2) Hydroboration Oxidation Reaction :



78. On reaction of Lead Sulphide with dilute nitric acid which of the following is **not** formed ?

- (1) Lead nitrate                      (2) Sulphur  
(3) Nitric oxide                      (4) Nitrous oxide

Ans. (4)



Nitrous oxide ( $\text{N}_2\text{O}$ ) is not formed during the reaction.

79. Identify the **incorrect** statements regarding primary standard of titrimetric analysis

- (A) It should be purely available in dry form.  
(B) It should not undergo chemical change in air.  
(C) It should be hygroscopic and should react with another chemical instantaneously and stoichiometrically.  
(D) It should be readily soluble in water.  
(E)  $\text{KMnO}_4$  &  $\text{NaOH}$  can be used as primary standard.

Choose the **correct** answer from the options given below :


- (1) (C) and (D) only                      (2) (B) and (E) only  
(3) (A) and (B) only                      (4) (C) and (E) only

Ans. (4)

Sol.  $\text{KMnO}_4$  &  $\text{NaOH} \rightarrow$  Secondary standard.

Primary standard should not be Hygroscopic.

80. 0.05M  $\text{CuSO}_4$  when treated with 0.01M  $\text{K}_2\text{Cr}_2\text{O}_7$  gives green colour solution of  $\text{Cu}_2\text{Cr}_2\text{O}_7$ . The [SPM : Semi Permeable Membrane]




Side X    SPM    Side Y

Due to osmosis :

- (1) Green colour formation observed on side Y.  
(2) Green colour formation observed on side X.  
(3) Molarity of  $\text{K}_2\text{Cr}_2\text{O}_7$  solution is lowered.  
(4) Molarity of  $\text{CuSO}_4$  solution is lowered.

Ans. (4)

Sol. Only solvent Molecules are allowed to pass through the SPM.




### SECTION-B

81. The heat of solution of anhydrous  $\text{CuSO}_4$  and  $\text{CuSO}_4 \cdot 5\text{H}_2\text{O}$  are  $-70 \text{ kJ mol}^{-1}$  and  $+12 \text{ kJ mol}^{-1}$  respectively.

The heat of hydration of  $\text{CuSO}_4$  to  $\text{CuSO}_4 \cdot 5\text{H}_2\text{O}$  is  $-x \text{ kJ}$ . The value of  $x$  is \_\_\_\_\_.

Ans. (82)



from (1) & (2)

$$-70 = x + 12$$

$$x = -82$$



Download the new **ALLEN** app  
& enroll for **Online Programs**


CLICK HERE TO  
DOWNLOAD



82. Given below are two statements :

**Statement I :** The rate law for the reaction  $A + B \rightarrow C$  is rate ( $r$ ) =  $k[A]^2[B]$ . When the concentration of both A and B is doubled, the reaction rate is increased "x" times.

**Statement II :**




The figure is showing "the variation in concentration against time plot" for a "y" order reaction.

The value of  $x + y$  is \_\_\_\_\_.


**Ans. (8)**

**Sol.**  $r = k[A]^2[B]$   
if conc. are doubled  
 $r' = k[2A]^2[2B]^1$   
 $r' = 8r \Rightarrow x = 8$




$\Rightarrow$  Zero order,  $y = 0$   
 $x + y = 8$




83. How many compounds among the following compounds show inductive, mesomeric as well as hyperconjugation effects?



**Ans. (4)**



84. The standard reduction potentials at 298 K for the following half cells are given below :



Consider the given electrochemical reactions, The number of metal(s) which will be oxidized by  $\text{Cr}_2\text{O}_7^{2-}$ , in aqueous solution is \_\_\_\_\_.

**Ans. (3)**

**Sol.** Fe, Ni, Ag will be oxidized due to lower S.R.P.

85. When equal volume of 1M HCl and 1M  $\text{H}_2\text{SO}_4$  are separately neutralised by excess volume of 1M NaOH solution. X and y kJ of heat is liberated respectively. The value of  $y/x$  is \_\_\_\_\_.

**Ans. (2)**

**Sol.**  $\text{H}^+ + \text{OH}^- \rightarrow \text{H}_2\text{O} \Rightarrow x$   
 $2\text{H}^+ + 2\text{OH}^- \rightarrow 2\text{H}_2\text{O} \Rightarrow 2x = y$   
 $y/x = 2$

86. Molarity (M) of an aqueous solution containing x g of anhyd.  $\text{CuSO}_4$  in 500 mL solution at  $32^\circ\text{C}$  is  $2 \times 10^{-1}$  M. Its molality will be \_\_\_\_\_  $\times 10^{-3}$  m. (nearest integer).

[Given density of the solution = 1.25 g/mL.]

**Allen Ans. (164)**

**NTA Ans. (81) BONUS**

**Sol.**  $M_{\text{sol}^n} = v_{\text{sol}^n} \times d_{\text{sol}^n}$   
 $= 500 \times 1.25 = 625\text{g}$

Mass of solute (x) =  $0.2 \times 0.5 \times 159.5$   
 $= 15.95$

$n_{\text{solute}} = 0.1,$

Mass of solvent = Mass of solution – Mass of solute  
 $= 625 - 15.95$   
 $= 609.05$

$m = \frac{0.1}{\frac{609.05}{1000}}$


$m = 0.164 = 164 \times 10^{-3}$



Download the new ALLEN app  
& enroll for Online Programs

CLICK HERE TO  
DOWNLOAD


87. The total number of species from the following in which one unpaired electron is present, is \_\_\_\_\_.



**Ans. (4)**

**Sol.** One unpaired  $e^-$  is present in :  $\text{C}_2^-$ ;  $\text{O}_2^-$ ;  $\text{H}_2^+$ ;  $\text{He}_2^+$

88. Number of ambidentate ligands among the following is \_\_\_\_\_.



**Ans. (3)**

**Sol.** Ligands which have two different donor sites but at a time connects with only one donor site to central metal are ambidentate ligands.

Ambidentate ligands are  $\text{NO}_2^-$ ;  $\text{SCN}^-$ ;  $\text{CN}^-$


89. Total number of essential amino acid among the given list of amino acids is \_\_\_\_\_.  
 Arginine, Phenylalanine, Aspartic acid, Cysteine, Histidine, Valine, Proline

**Ans. (4)**

**Sol.** Essential Amino acids are :-

Arginine, Phenylalanine, Histidine, Valine

90. Number of colourless lanthanoid ions among the following is \_\_\_\_\_.



**Ans. (2)**

**Sol.**  $\text{La}^{+3} - [\text{Xe}]4f^0$

$\text{Nd}^{+3} - [\text{Xe}]4f^3$

$\text{Sm}^{+3} - [\text{Xe}]4f^5$

$\text{Eu}^{+3} - [\text{Xe}]4f^6$

$\text{Lu}^{+3} - [\text{Xe}]4f^{14}$

$\text{La}^{+3}$  and  $\text{Lu}^{+3}$  do not show any colour because no unpaired electron is present.



Download the new **ALLEN** app  
& enroll for **Online Programs**

CLICK HERE TO  
DOWNLOAD

Are you targeting **JEE 2025** ?

Ace it with **ALLEN's**  
**Leader Course**

Online Program 18 APRIL '24

Offline Program 24 APRIL '24



**ALLEN**


Get The Latest  
**IIT-JEE** Special Books  
at Your Door Steps...!!

**JOIN THE JOURNEY OF LEARNING**

*with*

**SCORE TEST PAPERS | HANDBOOKS**  
**JEE-MAIN PYQ's | JEE-Adv. PYQ's**

**SHOP NOW**



Available in  
**HINDI & ENGLISH**