

## FINAL JEE-MAIN EXAMINATION – APRIL, 2024

(Held On Tuesday 09<sup>th</sup> April, 2024)

TIME : 3 : 00 PM to 6 : 00 PM

### PHYSICS

#### SECTION-A

31. A nucleus at rest disintegrates into two smaller nuclei with their masses in the ratio of 2:1. After disintegration they will move :-

- (1) In opposite directions with speed in the ratio of 1:2 respectively
- (2) In opposite directions with speed in the ratio of 2:1 respectively
- (3) In the same direction with same speed.
- (4) In opposite directions with the same speed.

**Ans. (1)**

**Sol.** By conservation of momentum


$$p_i = p_f$$

$$0 = m_1 u_1 + m_2 u_2$$

$$\frac{u_1}{u_2} = -\left[\frac{1}{2}\right] \text{ as } \frac{m_1}{m_2} = \frac{2}{1}$$

move in opposite direction with speed ratio 1 : 2


32. The following figure represents two biconvex lenses  $L_1$  and  $L_2$  having focal length 10 cm and 15 cm respectively. The distance between  $L_1$  &  $L_2$  is :



- (1) 10 cm
- (2) 15 cm
- (3) 25 cm
- (4) 35 cm

**Ans. (3)**

**Sol.**



$$D = f_1 + f_2 = 25 \text{ cm}$$

Paraxial parallel rays pass through focus and ray from focus of convex lens will become parallel

### TEST PAPER WITH SOLUTION

33. The temperature of a gas is  $-78^\circ\text{C}$  and the average translational kinetic energy of its molecules is  $K$ . The temperature at which the average translational kinetic energy of the molecules of the same gas becomes  $2K$  is :

- (1)  $-39^\circ\text{C}$
- (2)  $117^\circ\text{C}$
- (3)  $127^\circ\text{C}$
- (4)  $-78^\circ\text{C}$

**Ans. (2)**

**Sol.**  $K.E = \frac{nf_1RT}{2}$

$$T_i = -78^\circ\text{C} \rightarrow 273 + [-78^\circ\text{C}] = 195\text{K}$$

$$K.E \propto T$$

To double the K.E energy temp also become double

$$T_f = 390 \text{ K}$$

$$T_f = 117^\circ\text{C}$$

34. A hydrogen atom in ground state is given an energy of 10.2 eV. How many spectral lines will be emitted due to transition of electrons ?

- (1) 6
- (2) 3
- (3) 10
- (4) 1

**Ans. (4)**

**Sol.** Hydrogen will be in first excited state therefore it will emit one spectral line corresponding to transition b/w energy level 2 to 1

35. The magnetic field in a plane electromagnetic wave is  $B_y = (3.5 \times 10^{-7}) \sin(1.5 \times 10^3 x + 0.5 \times 10^{11} t)$  T. The corresponding electric field will be

- (1)  $E_y = 1.17 \sin(1.5 \times 10^3 x + 0.5 \times 10^{11} t) \text{Vm}^{-1}$
- (2)  $E_z = 105 \sin(1.5 \times 10^3 x + 0.5 \times 10^{11} t) \text{Vm}^{-1}$
- (3)  $E_z = 1.17 \sin(1.5 \times 10^3 x + 0.5 \times 10^{11} t) \text{Vm}^{-1}$
- (4)  $E_y = 10.5 \sin(1.5 \times 10^3 x + 0.5 \times 10^{11} t) \text{Vm}^{-1}$

**Ans. (2)**

**Sol.**  $E_0 = B_0 c$

$$E_0 = 3 \times 10^8 \times (3.5 \times 10^{-7}) \sin(1.5 \times 10^3 x + 0.5 \times 10^{11} t)$$

$$E_0 = 105 \sin(1.5 \times 10^3 x + 0.5 \times 10^{11} t) \text{Vm}^{-1}$$


Data inconsistent while calculating speed of wave. You can challenge for data.



Download the new **ALLEN app**  
& enroll for **Online Programs**

CLICK HERE TO  
DOWNLOAD


36. A square loop of side 15 cm being moved towards right at a constant speed of 2 cm/s as shown in figure. The front edge enters the 50 cm wide magnetic field at  $t = 0$ . The value of induced emf in the loop at  $t = 10$  s will be :



- (1) 0.3 mV                      (2) 4.5 mV  
(3) zero                         (4) 3 mV

Ans. (3)

Sol. At  $t = 10$  sec complete loop is in magnetic field therefore no change in flux




$$e = \frac{d\phi}{dt} = 0$$

$e = 0$  for complete loop

37. Two cars are travelling towards each other at speed of  $20 \text{ m s}^{-1}$  each. When the cars are 300 m apart, both the drivers apply brakes and the cars retard at the rate of  $2 \text{ m s}^{-2}$ . The distance between them when they come to rest is :

- (1) 200 m                      (2) 50 m  
(3) 100 m                      (4) 25 m

Ans. (3)



$$|\vec{u}_{BA}| = 40 \text{ m/s}$$

$$|\vec{a}_{BA}| = 4 \text{ m/s}^2$$


Apply  $(v^2 = u^2 + 2as)_{\text{relative}}$

$$0 = (40)^2 + 2(-4)(S)$$

$$S = 200 \text{ m}$$

$$\text{Remaining distance} = 300 - 200 = 100 \text{ m}$$

38. The  $I$ - $V$  characteristics of an electronic device shown in the figure. The device is :



- (1) a solar cell  
(2) a transistor which can be used as an amplifier  
(3) a zener diode which can be used as voltage regulator  
(4) a diode which can be used as a rectifier

Ans. (3)


Sol. Theory

Zener diode used as voltage regulator

39. The excess pressure inside a soap bubble is thrice the excess pressure inside a second soap bubble. The ratio between the volume of the first and the second bubble is :

- (1) 1 : 9                              (2) 1 : 3  
(3) 1 : 81                            (4) 1 : 27

Ans. (4)



Sol.

$$P_1 - P_0 = \frac{4T}{r_1}$$

$$P_2 - P_0 = \frac{4T}{r_2}$$

$$P_1 - P_0 = 3(P_2 - P_0)$$

$$\frac{4T}{r_1} = 3 \frac{4T}{r_2}$$

$$r_2 = 3r_1$$

$$\frac{V_1}{V_2} = \frac{\frac{4}{3}\pi r_1^3}{\frac{4}{3}\pi r_2^3} = \frac{1}{27}$$



Download the new ALLEN app  
& enroll for Online Programs

CLICK HERE TO  
DOWNLOAD


40. The de-Broglie wavelength associated with a particle of mass  $m$  and energy  $E$  is  $h/\sqrt{2mE}$ . The dimensional formula for Planck's constant is :
- (1)  $[ML^{-1}T^{-2}]$                       (2)  $[ML^2T^{-1}]$   
 (3)  $[MLT^{-2}]$                         (4)  $[M^2L^2T^{-2}]$

Ans. (2)

Sol.  $\lambda = \frac{h}{\sqrt{2mE}}$  or  $E = h\nu$   
 $[ML^2T^{-2}] = h[T^{-1}]$   
 $h = [ML^2T^{-1}]$

41. A satellite of  $10^3$  kg mass is revolving in circular orbit of radius  $2R$ . If  $\frac{10^4 R}{6}$  J energy is supplied to the satellite, it would revolve in a new circular orbit of radius :
- (use  $g = 10m/s^2$ ,  $R =$  radius of earth)
- (1)  $2.5 R$                                   (2)  $3 R$   
 (3)  $4 R$                                     (4)  $6 R$

Ans. (4)



Sol. Total energy =  $-\frac{GMm}{2(2R)}$

if energy =  $\frac{10^4 R}{6}$  is added then

$$\frac{-GMm}{4R} + \frac{10^4 R}{6} = \frac{-GMm}{2r}$$

where  $r$  is new radius of revolving and  $g = \frac{GM}{R^2}$


$$-\frac{mgR}{4} + \frac{10^4 R}{6} = -\frac{mgR^2}{2r} \quad (m = 10^3 \text{ kg})$$

$$-\frac{10^3 \times 10 \times R}{4} + \frac{10^4 R}{6} = -\frac{10^3 \times 10 \times R^2}{2r}$$

$$-\frac{1}{4} + \frac{1}{6} = -\frac{R}{2r}$$

$r = 6R$

42. The effective resistance between  $A$  and  $B$ , if resistance of each resistor is  $R$ , will be




- (1)  $\frac{2}{3} R$                                   (2)  $\frac{8R}{3}$   
 (3)  $\frac{5R}{3}$                                     (4)  $\frac{4R}{3}$


Ans. (2)

Sol. From symmetry we can remove two middle resistance.

New circuit is



43. Five charges  $+q$ ,  $+5q$ ,  $-2q$ ,  $+3q$  and  $-4q$  are situated as shown in the figure. The electric flux due to this configuration through the surface  $S$  is :



- (1)  $\frac{5q}{\epsilon_0}$                       (2)  $\frac{4q}{\epsilon_0}$   
 (3)  $\frac{3q}{\epsilon_0}$                       (4)  $\frac{q}{\epsilon_0}$

**Ans. (2)**

**Sol.** As per gauss theorem,

$$\phi = \frac{q_{in}}{\epsilon_0} = \frac{q + (-2q) + 5q}{\epsilon_0}$$

$$\frac{4q}{\epsilon_0}$$

44. A proton and a deuteron ( $q = +e$ ,  $m = 2.0u$ ) having same kinetic energies enter a region of uniform magnetic field  $\vec{B}$ , moving perpendicular to  $\vec{B}$ . The ratio of the radius  $r_d$  of deuteron path to the radius  $r_p$  of the proton path is :

- (1) 1 : 1                      (2) 1 :  $\sqrt{2}$   
 (3)  $\sqrt{2}$  : 1                      (4) 1 : 2

**Ans. (3)**

**Sol.** In uniform magnetic field,

$$R = \frac{mv}{qB} = \frac{\sqrt{2m(K.E)}}{qB}$$

Since same K.E

$$R \propto \frac{\sqrt{m}}{q}$$

$$\therefore \frac{R_{deuteron}}{R_{proton}} = \sqrt{\frac{m_d}{m_p}} \times \frac{q_p}{q_d}$$

$$= \sqrt{2} \times 1$$

$$\therefore \gamma_d : \gamma_p = \sqrt{2} : 1$$

45. UV light of 4.13 eV is incident on a photosensitive metal surface having work function 3.13 eV. The maximum kinetic energy of ejected photoelectrons will be :

- (1) 4.13 eV                      (2) 1 eV  
 (3) 3.13 eV                      (4) 7.26 eV

**Ans. (2)**

**Sol.**  $E_{\text{photon}} = (\text{work function}) + K.E_{\text{max}}$

$$\therefore 4.13 = 3.13 + K.E_{\text{max}}$$

$$\therefore K.E_{\text{max}} = 1 \text{ eV}$$

46. The energy released in the fusion of 2 kg of hydrogen deep in the sun is  $E_H$  and the energy released in the fission of 2 kg of  $^{235}\text{U}$  is  $E_U$ . The

ratio  $\frac{E_H}{E_U}$  is approximately :

(Consider the fusion reaction as  $4^1_1\text{H} + 2e^- \rightarrow ^4_2\text{He} + 2\nu + 6\gamma + 26.7 \text{ MeV}$ , energy

released in the fission reaction of  $^{235}\text{U}$  is 200 MeV per fission nucleus and  $N_A = 6.023 \times 10^{23}$ )

- (1) 9.13                      (2) 15.04  
 (3) 7.62                      (4) 25.6

**Ans. (3)**

**Sol.** In each fusion reaction, 4  $^1_1\text{H}$  nucleus are used.

Energy released per Nuclei of  $^1_1\text{H} = \frac{26.7}{4} \text{ MeV}$

$\therefore$  Energy released by 2 kg hydrogen ( $E_H$ )

$$= \frac{2000}{1} \times N_A \times \frac{26.7}{4} \text{ MeV}$$

&

$\therefore$  Energy released by 2 kg Uranium ( $E_U$ )

$$= \frac{2000}{235} \times N_A \times 200 \text{ MeV}$$

So,

$$\frac{E_H}{E_U} = 235 \times \frac{26.7}{4 \times 200} = 7.84$$


$\therefore$  Approximately close to 7.62



Download the new **ALLEN** app  
 & enroll for **Online Programs**

CLICK HERE TO  
 DOWNLOAD

47. A real gas within a closed chamber at  $27^\circ\text{C}$  undergoes the cyclic process as shown in figure. The gas obeys  $PV^3 = RT$  equation for the path  $A$  to  $B$ . The net work done in the complete cycle is (assuming  $R = 8\text{J/molK}$ ):



- (1) 225 J                                      (2) 205 J  
 (3) 20 J                                        (4) -20 J

Ans. (2)

Sol.  $W_{AB} = \int PdV$  (Assuming T to be constant)

$$= \int \frac{RTdV}{V^3}$$

$$= RT \int_2^4 V^{-3} dV$$

$$= 8 \times 300 \times \left( -\frac{1}{2} \left[ \frac{1}{4^2} - \frac{1}{2^2} \right] \right)$$

$$= 225 \text{ J}$$

$$W_{BC} = P \int_4^2 dV = 10(2 - 4) = -20 \text{ J}$$


$$W_{CA} = 0$$

$$\therefore W_{\text{cycle}} = 205 \text{ J}$$

Note : Data is inconsistent in process AB.

So needs to be challenged.

48. A 1 kg mass is suspended from the ceiling by a rope of length 4m. A horizontal force 'F' is applied at the mid point of the rope so that the rope makes an angle of  $45^\circ$  with respect to the vertical axis as shown in figure. The magnitude of F is :



- (1)  $\frac{10}{\sqrt{2}} \text{ N}$                                       (2) 1 N  
 (3)  $\frac{1}{10 \times \sqrt{2}} \text{ N}$                                       (4) 10 N

Ans. (4)

Sol.  $T_1 \sin 45^\circ = F$

$$T_1 \cos 45^\circ = T_2 = 1 \times g$$

$$\therefore \tan 45^\circ = \frac{F}{g}$$

$$\therefore F = 10 \text{ N}$$

49. A spherical ball of radius  $1 \times 10^{-4} \text{ m}$  and density  $10^5 \text{ kg/m}^3$  falls freely under gravity through a distance  $h$  before entering a tank of water, If after entering in water the velocity of the ball does not change, then the value of  $h$  is approximately :

(The coefficient of viscosity of water is  $9.8 \times 10^{-6} \text{ N s/m}^2$ )

- (1) 2296 m                                      (2) 2249 m  
 (3) 2518 m                                      (4) 2396 m

Ans. (3)

Sol.  $V_T = \frac{2g R^2 [\rho_B - \rho_L]}{9 \eta}$


$\Rightarrow V_T = \frac{2}{9} \times \frac{10 \times (10^{-4})^2}{9.8 \times 10^{-6}} [10^5 - 10^3]$

$\Rightarrow V_T = 224.5$

when ball fall from height (h)

$V = \sqrt{2gh}$

$h = \left(\frac{V^2}{2g}\right) = 2518\text{m}$




In the truth table of the above circuit the value of X and Y are :


- (1) 1, 1                      (2) 1, 0  
(3) 0, 1                      (4) 0, 0

Ans. (1)

Sol. For x



For y




SECTION-B

51. A straight magnetic strip has a magnetic moment of  $44 \text{ Am}^2$ . If the strip is bent in a semicircular shape, its magnetic moment will be .....  $\text{Am}^2$   
(Given  $\pi = \frac{22}{7}$ )

Ans. (28)

Sol. Magnetic moment of straight wire =  $m \times \ell = 44$



Magnetic moment of arc

$= m \times 2r$

$= m \times \frac{2\ell}{\pi}$

$= \frac{44 \times 2}{\pi} = \frac{88}{\pi} = 28$

52. A particle of mass 0.50 kg executes simple harmonic motion under force  $F = -50(\text{Nm}^{-1})x$ . The time period of oscillation is  $\frac{x}{35}$ s. The value of x is

.....

(Given  $\pi = \frac{22}{7}$ )

Ans. (22)

Sol.  $m = 0.5 \text{ kg}$

$F = -50(x)$

$ma = (-50x)$


$0.5 a = -50x$

$a = (-100x)$

$\omega^2 = 100 \Rightarrow (\omega = 10)$

$T = \frac{2\pi}{10} = \left(\frac{\pi}{5}\right) = \frac{22}{7 \times 5} = \left(\frac{22}{35}\right)$

$\frac{\pi}{35} = \frac{22}{35} \Rightarrow \boxed{x = 22}$




Download the new ALLEN app & enroll for Online Programs

CLICK HERE TO DOWNLOAD



53. A capacitor of reactance  $4\sqrt{3}\Omega$  and a resistor of resistance  $4\Omega$  are connected in series with an ac source of peak value  $8\sqrt{2}V$ . The power dissipation in the circuit is .....W.

Ans. (4)



$$Z = \sqrt{R^2 + X^2}$$


$$Z = \sqrt{4^2 + (4\sqrt{3})^2} = 8\Omega$$

$$V_{rms} = \frac{V}{\sqrt{2}} = \frac{8\sqrt{2}}{\sqrt{2}} = (8V)$$


$$I_{rms} = \frac{V_{rms}}{Z} = \frac{8}{8} = 1A$$

$$\text{Power dissipated} = I_{rms}^2 \times R = 1 \times 4 = (4W)$$

54. An electric field  $\vec{E} = (2x\hat{i})NC^{-1}$  exists in space. A cube of side 2m is placed in the space as per figure given below. The electric flux through the cube is .....  $Nm^2/C$ .



Ans. (16)



$$\phi_{in} = -4 \times 4 = -16 Nm^2 / c$$

$$\phi_{out} = 8 \times 4 = 32 Nm^2 / c$$

$$\phi_{net} = \phi_{in} + \phi_{out} = -16 + 32 = 16 Nm^2 / c$$

55. A circular disc reaches from top to bottom of an inclined plane of length  $l$ . When it slips down the plane, it takes  $t$  s. When it rolls down the plane then it takes  $\left(\frac{\alpha}{2}\right)^{1/2} t$  s, where  $\alpha$  is .....

Ans. (3)

Sol. For slipping

$$a = g \sin \theta$$

$$l = \frac{1}{2} at^2 \Rightarrow t = \sqrt{\frac{2l}{g \sin \theta}}$$

For rolling

$$a' = \frac{g \sin \theta}{1 + \frac{k^2}{R^2}} \left[ k = \frac{R}{\sqrt{2}} \right]$$


$$\Rightarrow a' = \frac{2g \sin \theta}{3}$$


$$l = \frac{1}{2} a' (t')^2$$

$$\Rightarrow t' = \sqrt{\frac{6l}{2g \sin \theta}} = \sqrt{\frac{\alpha}{2}} \sqrt{\frac{2l}{g \sin \theta}}$$

$$\Rightarrow \alpha = 3$$

56. To determine the resistance (R) of a wire, a circuit is designed below, The V-I characteristic curve for this circuit is plotted for the voltmeter and the ammeter readings as shown in figure. The value of R is .....  $\Omega$ .





Ans. (2500)

Sol.  $R_{eq} = \frac{10^4 R}{10^4 + R}$

$E = 4V, I = 2mA$

$I = \frac{E}{R_{eq}} \Rightarrow 2 \times 10^{-3} = \frac{4(10^4 + R)}{10^4 R}$


$\Rightarrow 20R = 40000 + 4R$

$16R = 40000$

$R = 2500\Omega$

57. The resultant of two vectors  $\vec{A}$  and  $\vec{B}$  is perpendicular to  $\vec{A}$  and its magnitude is half that of  $\vec{B}$ . The angle between vectors  $\vec{A}$  and  $\vec{B}$  is .....

Ans. (150)



$B \cos \theta = \frac{B}{2}$

$\Rightarrow \theta = 60^\circ$

So, angle between  $\vec{A}$  &  $\vec{B}$  is  $90^\circ + 60^\circ = 150^\circ$

58. Monochromatic light of wavelength 500 nm is used in Young's double slit experiment. An interference pattern is obtained on a screen. When one of the slits is covered with a very thin glass plate (refractive index = 1.5), the central maximum is shifted to a position previously occupied by the 4<sup>th</sup> bright fringe. The thickness of the glass-plate is ..... $\mu\text{m}$ .

Ans. (4)

Sol.  $(\mu - 1)t = n\lambda$

$(1.5 - 1)t = 4 \times 500 \times 10^{-9} \text{ m}$

$t = 4000 \times 10^{-9} \text{ m}$

$t = 4\mu\text{m}$

59. A force  $(3x^2 + 2x - 5)$  N displaces a body from  $x = 2$  m to  $x = 4$  m. Work done by this force is .....J.

Ans. (58)

Sol.  $W = \int_{x_1}^{x_2} F dx$

$W = \int_2^4 (3x^2 + 2x - 5) dx$

$W = [x^3 + x^2 - 5x]_2^4$

$W = [60 - 2]J = 58J$

60. At room temperature ( $27^\circ\text{C}$ ), the resistance of a heating element is  $50\Omega$ . The temperature coefficient of the material is  $2.4 \times 10^{-4} \text{ }^\circ\text{C}^{-1}$ . The temperature of the element, when its resistance is  $62\Omega$ , is ..... $^\circ\text{C}$ .

Ans. (1027)

Sol.  $R = R_0(1 + \alpha\Delta T)$

$62 = 50 [1 + 2.4 \times 10^{-4} \Delta T]$

$\Delta T = 1000^\circ\text{C}$

$\Rightarrow T - 27^\circ = 1000^\circ\text{C}$

$T = 1027^\circ\text{C}$



Download the new **ALLEN** app  
& enroll for **Online Programs**

CLICK HERE TO  
DOWNLOAD





# Are you targeting **JEE 2025** ?

## Ace it with **ALLEN's** **Leader Course**

Online Program 18 APRIL '24

Offline Program 24 APRIL '24



# ALLEN


Get The Latest  
**IIT-JEE** Special Books  
at Your Door Steps...!!

**JOIN THE JOURNEY OF LEARNING**

*with*

**SCORE TEST PAPERS | HANDBOOKS**  
**JEE-MAIN PYQ's | JEE-Adv. PYQ's**

**SHOP NOW**



Available in  
**HINDI & ENGLISH**