

FINAL JEE-MAIN EXAMINATION – JANUARY, 2023

(Held On Monday 30th January, 2023)

TIME : 9 : 00 AM to 12 : 00 NOON

MATHEMATICS

SECTION-A

61. Let $A = \begin{bmatrix} m & n \\ p & q \end{bmatrix}$, $d = |A| \neq 0$ $|A - d(\text{adj } A)| = 0$. Then

(1) $(1+d)^2 = (m+q)^2$

(2) $1+d^2 = (m+q)^2$

(3) $(1+d)^2 = m^2 + q^2$

(4) $1+d^2 = m^2 + q^2$

Official Ans. by NTA (1)

Allen Ans. (1)

Sol. $A = \begin{bmatrix} m & n \\ p & q \end{bmatrix}$, $|A - d(\text{adj } A)| = 0$

$$\Rightarrow |A - d(\text{adj } A)| = \left| \begin{bmatrix} m & n \\ p & q \end{bmatrix} - d \begin{bmatrix} q & -n \\ -p & m \end{bmatrix} \right| = \left| \begin{matrix} m - qd & n(1+d) \\ p(1+d) & q - md \end{matrix} \right| = 0$$

$$\Rightarrow (m - qd)(q - md) - np(1 + d)^2 = 0$$

$$\Rightarrow mq - m^2d - q^2d + mqd^2 - np(1 + d)^2 = 0$$

$$\Rightarrow (mq - np) + d^2(mq - np) - d(m^2 + q^2 + 2np) = 0$$

$$\Rightarrow d + d^3 - d((m+q)^2 - 2d) = 0$$

$$\Rightarrow 1 + d^2 = (m+q)^2 - 2d$$

$$\Rightarrow (1+d)^2 = (m+q)^2$$

∴ Option (1) is correct.

62. The line l_1 passes through the point $(2,6,2)$ and is perpendicular to the plane $2x + y - 2z = 10$. Then the shortest distance between the line l_1 and the line

$$\frac{x+1}{2} = \frac{y+4}{-3} = \frac{z}{2} \text{ is :}$$

(1) 7

(2) $\frac{19}{3}$

(3) $\frac{19}{3}$

(4) 9

Official Ans. by NTA (4)

Allen Ans. (4)


TEST PAPER WITH SOLUTION

Sol. Line ℓ , is given by

$$L_1: \frac{x-2}{2} = \frac{y-6}{1} = \frac{z-2}{-2}$$

Given,

$$L_2: \frac{x+1}{2} = \frac{y+4}{-3} = \frac{z}{2}$$



$$\text{Shortest distance} = \frac{|\overrightarrow{AB} \cdot \overrightarrow{MN}|}{\overrightarrow{MN}}$$

$$\overrightarrow{AB} = 3\hat{i} + 10\hat{j} + 2\hat{k}$$

$$\overrightarrow{MN} = \begin{vmatrix} \hat{i} & \hat{j} & \hat{k} \\ 2 & 1 & -2 \\ 2 & -3 & 2 \end{vmatrix} = -4\hat{i} - 8\hat{j} - 8\hat{k}$$

$$MN = \sqrt{16 + 64 + 64} = 12$$

$$\therefore \text{Shortest distance} = \frac{|-12 - 80 - 16|}{12} = 9$$

∴ Option (4) is correct.

63. If an unbiased die, marked with $-2, -1, 0, 1, 2, 3$ on its faces, is thrown five times, then the probability that the product of the outcomes is positive, is :

(1) $\frac{881}{2592}$

(2) $\frac{521}{2592}$

(3) $\frac{440}{2592}$

(4) $\frac{27}{288}$

Official Ans. by NTA (2)

Allen Ans. (2)

Sol. Either all outcomes are positive or any two are negative.

$$\text{Now, } p = P(\text{positive}) = \frac{3}{6} = \frac{1}{2}$$

$$q = p(\text{negative}) = \frac{2}{6} = \frac{1}{3}$$

Required probability

$$= {}^5C_5 \left(\frac{1}{2}\right)^5 + {}^5C_2 \left(\frac{1}{3}\right)^2 \left(\frac{1}{2}\right)^3 + {}^5C_4 \left(\frac{1}{3}\right)^4 \left(\frac{1}{2}\right)^1 \\ = \frac{521}{2592}$$

\therefore Option (2) is correct.

64. Let the system of linear equations

$$x + y + kz = 2$$

$$2x + 3y - z = 1$$

$$3x + 4y + 2z = k$$

have infinitely many solutions. Then the system

$$(k+1)x + (2k-1)y = 7$$

$$(2k+1)x + (k+5)y = 10 \text{ has :}$$

(1) infinitely many solutions

(2) unique solution satisfying $x - y = 1$

(3) no solution

(4) unique solution satisfying $x + y = 1$

Official Ans. by NTA (4)

Allen Ans. (4)

$$\begin{vmatrix} 1 & 1 & k \\ 2 & 3 & -1 \\ 3 & 4 & 2 \end{vmatrix} = 0$$

$$\Rightarrow 1(10) - 1(7) + k(-1) = 0$$

$$\Rightarrow k = 3$$

For $k = 3$, 2nd system is

$$4x + 5y = 7 \quad \dots\dots(1)$$

$$\text{and } 7x + 8y = 10 \quad \dots\dots(2)$$

Clearly, they have a unique solution

$$(2) - (1) \Rightarrow 3x + 3y = 3$$

$$\Rightarrow x + y = 1$$

65. If $\tan 15^\circ + \frac{1}{\tan 75^\circ} + \frac{1}{\tan 105^\circ} + \tan 195^\circ = 2a$,
then the value of $\left(a + \frac{1}{a}\right)$ is :

$$(1) 4 \quad (2) 4 - 2\sqrt{3}$$

$$(3) 2 \quad (4) 5 - \frac{3}{2}\sqrt{3}$$

Official Ans. by NTA (1)

Allen Ans. (1)

Sol. Option (1)

$$\tan 15^\circ = 2 - \sqrt{3}$$

$$\frac{1}{\tan 75^\circ} = \cot 75^\circ = 2 - \sqrt{3}$$

$$\frac{1}{\tan 105^\circ} = \cot(105^\circ) = -\cot 75^\circ = \sqrt{3} - 2$$

$$\tan 195^\circ = \tan 15^\circ = 2 - \sqrt{3}$$

$$\therefore 2(2 - \sqrt{3}) = 2a \Rightarrow a = 2 - \sqrt{3}$$

$$\Rightarrow a + \frac{1}{a} = 4$$

66. Suppose $f : \mathbb{R} \rightarrow (0, \infty)$ be a differentiable function such that $5f(x+y) = f(x) \cdot f(y), \forall x, y \in \mathbb{R}$. If

$$f(3) = 320, \text{ then } \sum_{n=0}^5 f(n) \text{ is equal to :}$$

$$(1) 6875 \quad (2) 6575$$

$$(3) 6825 \quad (4) 6528$$

Official Ans. by NTA (3)

Allen Ans. (3)

Sol. Option (3)

$$5f(x+y) = f(x) \cdot f(y)$$

$$5f(0) = f(0)^2 \Rightarrow f(0) = 5$$

$$5f(x+1) = f(x) \cdot f(1)$$

$$\Rightarrow \frac{f(x+1)}{f(x)} = \frac{f(1)}{5}$$

$$\Rightarrow \frac{f(1)}{f(0)} \cdot \frac{f(2)}{f(1)} \cdot \frac{f(3)}{f(2)} = \left(\frac{f(1)}{5}\right)^3$$

$$\Rightarrow \frac{320}{5} = \frac{(f(1))^3}{5^3} \Rightarrow f(1) = 20$$

$$\therefore 5f(x+1) = 20 \cdot f(x) \Rightarrow f(x+1) = 4f(x)$$

$$\sum_{n=0}^5 f(n) = 5 + 5 \cdot 4 + 5 \cdot 4^2 + 5 \cdot 4^3 + 5 \cdot 4^4 + 5 \cdot 4^5$$

$$= \frac{5[4^6 - 1]}{3} = 6825$$

67. If $a_n = \frac{-2}{4n^2 - 16n + 15}$, then $a_1 + a_2 + \dots + a_{25}$ is equal to :

- (1) $\frac{51}{144}$ (2) $\frac{49}{138}$
 (3) $\frac{50}{141}$ (4) $\frac{52}{147}$

Official Ans. by NTA (3)

Allen Ans. (3)

Sol. Option (3)

$$\text{If } a_n = \frac{-2}{4n^2 - 16n + 15} \text{ then } a_1 + a_2 + \dots + a_{25}$$

$$\Rightarrow \sum_{n=1}^{25} a_n = \sum \frac{-2}{4n^2 - 16n + 15}$$

$$= \sum \frac{-2}{4n^2 - 6n - 10n + 15}$$

$$= \sum \frac{-2}{2n(2n-3) - 5(2n-3)}$$

$$= \sum \frac{-2}{(2n-3)(2n-5)}$$

$$= \sum \frac{1}{2n-3} - \frac{1}{2n-5}$$

$$= \frac{1}{47} - \frac{1}{(-3)}$$

$$= \frac{50}{141}$$

68. If the coefficient of x^{15} in the expansion of $\left(ax^3 + \frac{1}{bx^3}\right)^{15}$ is equal to the coefficient of x^{-15} in

the expansion of $\left(ax^{\frac{1}{3}} - \frac{1}{bx^3}\right)^{15}$, where a and b

are positive real numbers, then for each such ordered pair (a, b) :

- (1) $a = b$ (2) $ab = 1$
 (3) $a = 3b$ (4) $ab = 3$

Official Ans. by NTA (2)

Allen Ans. (2)

Sol. Option (2)

Coefficient Of x^{15} in $\left(ax^3 + \frac{1}{bx^3}\right)^{15}$

$$T_{r+1} = {}^{15}C_r \left(ax^3\right)^{15-r} \left(\frac{1}{bx^3}\right)^r$$

$$45 - 3r - \frac{r}{3} = 15$$

$$30 = \frac{10r}{3}$$

$$r = 9$$

Coefficient of $x^{15} = {}^{15}C_9 a^6 b^{-9}$

Coefficient of x^{-15} in $\left(ax^{\frac{1}{3}} - \frac{1}{bx^3}\right)^{15}$

$$T_{r+1} = {}^{15}C_r \left(ax^{\frac{1}{3}}\right)^{15-r} \left(-\frac{1}{bx^3}\right)^r$$

$$5 - \frac{r}{3} - 3r = -15$$

$$\frac{10r}{3} = 20$$

$$r = 6$$

Coefficient = ${}^{15}C_6 a^9 \times b^{-6}$

$$\Rightarrow \frac{a^9}{b^6} = \frac{a^6}{b^9}$$

$$\Rightarrow a^3 b^3 = 1 \Rightarrow ab = 1$$

69. If $\vec{a}, \vec{b}, \vec{c}$ are three non-zero vectors and \hat{n} is a unit vector perpendicular to \vec{c} such that $\vec{a} = \alpha \vec{b} - \hat{n}$, ($\alpha \neq 0$) and $\vec{b} \cdot \vec{c} = 12$, then

$|\vec{c} \times (\vec{a} \times \vec{b})|$ is equal to :

- (1) 15
 (2) 9
 (3) 12
 (4) 6

Official Ans. by NTA (3)

Allen Ans. (3)

73. Let a unit vector \widehat{OP} make angle α, β, γ with the positive directions of the co-ordinate axes OX, OY, OZ respectively, where $\beta \in \left(0, \frac{\pi}{2}\right)$ \widehat{OP} is perpendicular to the plane through points (1, 2, 3), (2, 3, 4) and (1, 5, 7), then which one of the following is true?

- (1) $\alpha \in \left(\frac{\pi}{2}, \pi\right)$ and $\gamma \in \left(\frac{\pi}{2}, \pi\right)$
- (2) $\alpha \in \left(0, \frac{\pi}{2}\right)$ and $\gamma \in \left(0, \frac{\pi}{2}\right)$
- (3) $\alpha \in \left(\frac{\pi}{2}, \pi\right)$ and $\gamma \in \left(0, \frac{\pi}{2}\right)$
- (4) $\alpha \in \left(0, \frac{\pi}{2}\right)$ and $\gamma \in \left(\frac{\pi}{2}, \pi\right)$

Official Ans. by NTA (1)

Allen Ans. (1)

Sol. Equation of plane :-

$$\begin{vmatrix} x-1 & y-2 & z-3 \\ 1 & 1 & 1 \\ 0 & 3 & 4 \end{vmatrix} = 0$$

$$\Rightarrow [x-1] - 4[y-2] + 3[z-3] = 0$$

$$\Rightarrow x - 4y + 3z = 2$$

D.R's of normal of plane $\langle 1, -4, 3 \rangle$

D.C's of $\left\langle \pm \frac{1}{\sqrt{26}}, \mp \frac{4}{\sqrt{26}}, \pm \frac{3}{\sqrt{26}} \right\rangle$

$$\cos \beta = \frac{4}{\sqrt{26}}$$

$$\cos \alpha = \frac{-1}{\sqrt{26}}$$

$$\cos \gamma = \frac{-3}{\sqrt{26}}$$

Ans. : (1)

74. If [t] denotes the greatest integer ≤ 1 , then the value

of $\frac{3(e-1)^2}{e} \int_1^2 x^2 e^{[x]+[x^3]} dx$ is :

- (1) $e^9 - e$
- (2) $e^8 - e$
- (3) $e^7 - 1$
- (4) $e^8 - 1$

Official Ans. by NTA (2)

Allen Ans. (2)

Sol.
$$\int_1^2 x^2 e^{[x^3]+1} dx$$

$$x^3 = t$$

$$3x^2 dx = dt$$

$$= \frac{e}{3} \int_1^8 e^{[t]} dt$$

$$= \frac{e}{3} \left\{ \int_1^2 e^{dt} + \int_2^3 e^{2t} dt + \dots + \int_7^8 e^{7t} dt \right\}$$

$$= \frac{e}{3} (e + e^2 + \dots + e^7)$$

$$= \frac{e^2}{3} (1 + e + \dots + e^6) = \frac{e^2}{3} \frac{(e^7 - 1)}{(e - 1)}$$

$$\frac{3(e-1)}{e} \int_1^2 x^2 \times e^{[x]+[x^3]} dx = \frac{3}{e} (e-1) \times \frac{e^2}{3} \frac{(e^7 - 1)}{(e - 1)}$$

$$= e(e^7 - 1)$$

$$= e^8 - e$$

Ans. : (2)

75. If P(h,k) be point on the parabola $x = 4y^2$, which is nearest to the point Q(0,33), then the distance of P from the directrix of the parabola $y^2 = 4(x+y)$ is equal to :

- (1) 2
- (2) 4
- (3) 8
- (4) 6

Official Ans. by NTA (4)

Allen Ans. (4)

Sol. Equation of normal

$$y = -tx + 2at + at^3$$

$$y = -tx + \frac{2}{16}t + \frac{1}{16}t^3$$

It passes through (0, 33)

$$33 = \frac{t}{8} + \frac{t^3}{16}$$

$$t^3 + 2t - 528 = 0$$

$$(t-8)(t^2 + 8t + 66) = 0$$

$$t = 8$$

$$P(at^2, 2at) = \left(\frac{1}{16} \times 64, 2 \times \frac{1}{16} \times 8 \right) = (4, 1)$$

Parabola :

$$y^2 = 4(x+y)$$

$$\Rightarrow y^2 - 4y = 4x$$

$$\Rightarrow (y-2)^2 = 4(x+1)$$

Equation of directix :-


$$x + 1 = -1$$

$$x = -2$$

Distance of point = 6

Ans. : (4)

76. A straight line cuts off the intercepts $OA = a$ and $OB = b$ on the positive directions of x -axis and y -axis respectively. If the perpendicular from origin O to this line makes an angle of $\frac{\pi}{6}$ with positive direction of y -axis and the area of ΔOAB is $\frac{98}{3}\sqrt{3}$, then $a^2 - b^2$ is equal to:
- (1) $\frac{392}{3}$ (2) 196
(3) $\frac{196}{3}$ (4) 98

Official Ans. by NTA (1)
Allen Ans. (1)
Sol.


$$\text{Equation of straight line : } \frac{x}{a} + \frac{y}{b} = 1$$

$$\text{Or } x \cos \frac{\pi}{3} + y \sin \frac{\pi}{3} = p$$

$$\frac{x}{2} + \frac{y\sqrt{3}}{2} = p$$

$$\frac{x}{3p} + \frac{y}{2p} = 1$$

$$\text{Comparing both : } a = 2p, b = \frac{2p}{\sqrt{3}}$$

$$\text{Now area of } \Delta OAB = \frac{1}{2} \cdot ab = \frac{98}{3} \cdot \sqrt{3}$$

$$\frac{1}{2} \cdot 2p \cdot \frac{2p}{\sqrt{3}} = \frac{98}{3} \cdot \sqrt{3}$$

$$p^2 = 49$$

$$a^2 - b^2 = 4p^2 - \frac{4p^2}{3} = \frac{2}{3}4p^2$$

$$= \frac{8}{3} \cdot 49 = \frac{392}{3}$$

77. The coefficient of x^{301} in $(1+x)^{500} + x(1+x)^{499} + x^2(1+x)^{498} + \dots + x^{500}$ is:
- (1) ${}^{501}C_{302}$ (2) ${}^{500}C_{301}$
(3) ${}^{500}C_{300}$ (4) ${}^{501}C_{200}$

Official Ans. by NTA (4)
Allen Ans. (4)

$$\text{Sol. } (1+x)^{500} + x(1+x)^{499} + x^2(1+x)^{498} + \dots + x^{500}$$

$$= (1+x)^{500} \cdot \left\{ \frac{1 - \left(\frac{x}{1+x} \right)^{501}}{1 - \frac{x}{1+x}} \right\}$$

$$= (1+x)^{500} \frac{\left((1+x)^{501} - x^{501} \right)}{(1+x)^{501}} \cdot (1+x)$$

$$= (1+x)^{501} - x^{501}$$

Coefficient of x^{301} in $(1+x)^{501} - x^{501}$ is given by

$${}^{501}C_{301} = {}^{501}C_{200}$$

78. Among the statements:

$$(S1) \quad ((p \vee q) \Rightarrow r) \Leftrightarrow (p \Rightarrow r)$$

$$(S2) \quad ((p \vee q) \Rightarrow r) \Leftrightarrow ((p \Rightarrow r) \vee (q \Rightarrow r))$$

(1) Only (S1) is a tautology

(2) Neither (S1) nor (S2) is a tautology

(3) Only (S2) is a tautology

(4) Both (S1) and (S2) are tautologies

Official Ans. by NTA (2)
Allen Ans. (2)

$$\text{Sol. } S_1 \equiv ((p \vee q) \Rightarrow r) \Leftrightarrow (p \Rightarrow r)$$

$$\begin{array}{ccccccccc} p & q & r & p \vee q & (p \vee q) \Rightarrow r & p \Rightarrow r & ((p \vee q) \Rightarrow r) \Leftrightarrow (p \Rightarrow r) \\ \hline T & T & T & T & T & T & T & T \\ T & T & F & T & F & F & F & T \\ T & F & T & T & T & T & T & T \\ F & T & T & T & T & T & T & T \\ T & F & F & T & F & F & F & T \\ F & T & F & T & F & T & T & T \\ F & F & F & T & T & T & T & T \\ F & F & F & F & T & T & T & T \end{array}$$

case – 1

$f(6) = S$ i.e. 1 option,

$f(5) = \text{any 5 element subset A of } S \text{ i.e. 6 options,}$

$f(4) = \text{any 4 element subset B of A i.e. 5 options,}$

$f(3) = \text{any 3 element subset C of B i.e. 4 options,}$

$f(2) = \text{any 2 element subset D of C i.e. 3 options,}$

$f(1) = \text{any 1 element subset E of D or empty subset i.e. 3 options,}$

Total functions = 1080

Case – 2

$f(6) = \text{any 5 element subset A of } S \text{ i.e. 6 options,}$

$f(5) = \text{any 4 element subset B of A i.e. 5 options,}$

$f(4) = \text{any 3 element subset C of B i.e. 4 options,}$

$f(3) = \text{any 2 element subset D of C i.e. 3 options,}$

$f(2) = \text{any 1 element subset E of D i.e. 2 options,}$

$f(1) = \text{empty subset i.e. 1 option}$

Total functions = 720

Case – 3

$f(6) = S$

$f(5) = \text{any 4 element subset A of } S \text{ i.e. 15 options,}$

$f(4) = \text{any 3 element subset B of A i.e. 4 options,}$

$f(3) = \text{any 2 element subset C of B i.e. 3 options,}$

$f(2) = \text{any 1 element subset D of C i.e. 2 options,}$

$f(1) = \text{empty subset i.e. 1 option}$

Total functions = 360

Case – 4

$f(6) = S$

$f(5) = \text{any 5 element subset A of } S \text{ i.e. 6 options,}$

$f(4) = \text{any 4 element subset B of A i.e. 10 options,}$

$f(3) = \text{any 3 element subset C of B i.e. 10 options,}$

$f(2) = \text{any 2 element subset D of C i.e. 2 options,}$

$f(1) = \text{empty subset i.e. 1 option}$

Total functions = 360

Case – 5

$f(6) = S$

$f(5) = \text{any 5 element subset A of } S \text{ i.e. 6 options,}$

$f(4) = \text{any 4 element subset B of A i.e. 5 options,}$

$f(3) = \text{any 3 element subset C of B i.e. 6 options,}$

$f(2) = \text{any 2 element subset D of C i.e. 2 options,}$

$f(1) = \text{empty subset i.e. 1 option}$

Total functions = 360

Case – 6

$f(6) = S$

$f(5) = \text{any 5 element subset A of } S \text{ i.e. 6 options,}$

$f(4) = \text{any 4 element subset B of A i.e. 5 options,}$

$f(3) = \text{any 3 element subset C of B i.e. 4 options,}$

$f(2) = \text{any 2 element subset D of C i.e. 3 options,}$

$f(1) = \text{empty subset i.e. 1 option}$

Total functions = 360


\therefore Number of such functions = 3240

82. Let α be the area of the larger region bounded by the curve $y^2 = 8x$ and the lines $y = x$ and $x = 2$, which lies in the first quadrant. Then the value of 3α is equal to _____.

Official Ans. by NTA (22)

Allen Ans. (22)

Sol.



Solving it

$$x^2 = 8x$$

$$\therefore x = 0, 8$$

$$\therefore y = 0, 8$$

$x = 2$ will intersect occur at

$$y^2 = 16 \Rightarrow y = \pm 4$$

\therefore Area of shaded

$$= \int_2^8 (\sqrt{8x} - x) dx = \int_2^8 (2\sqrt{2}\sqrt{x} - x) dx$$

$$= \left[2\sqrt{2} \cdot \frac{x^{3/2}}{3/2} - \frac{x^2}{2} \right]_0^8$$

$$= \left(\frac{4\sqrt{2}}{3} \cdot 2^{9/2} - 32 \right) - \left(\frac{4\sqrt{2}}{3} \cdot 2^{9/2} - 2 \right)$$

$$= \frac{128}{3} - 32 - \frac{16}{3} + 2 = \frac{112 - 90}{3} = \frac{22}{3} = A$$

$$\therefore 3A = 22$$

83. If $\lambda_1 < \lambda_2$ are two values of λ such that the angle between the planes $P_1 : \vec{r} \cdot (3\hat{i} - 5\hat{j} + \hat{k}) = 7$ and $P_2 : \vec{r} \cdot (\lambda\hat{i} + \hat{j} - 3\hat{k}) = 9$ is $\sin^{-1}\left(\frac{2\sqrt{6}}{5}\right)$, then the square of the length of perpendicular from the point $(38\lambda_1, 10\lambda_2, 2)$ to the plane P_1 is ____.

Official Ans. by NTA (315)

Allen Ans. (315)

$$\text{Sol. } P_1 = \vec{r} \cdot (3\hat{i} - 5\hat{j} + \hat{k}) = 7$$

$$P_2 = \vec{r} \cdot (\lambda\hat{i} + \hat{j} - 3\hat{k}) = 9$$

$$\theta = \sin^{-1}\left(\frac{2\sqrt{6}}{5}\right)$$

$$\Rightarrow \sin \theta = \frac{2\sqrt{6}}{5}$$

$$\therefore \cos \theta = \frac{1}{5}.$$

$$\cos \theta = \frac{\vec{r}_1 \cdot \vec{r}_2}{|\vec{r}_1| |\vec{r}_2|}$$

$$= \frac{(3\hat{i} - 5\hat{j} + \hat{k})(\lambda\hat{i} + \hat{j} - 3\hat{k})}{\sqrt{35} \cdot \sqrt{\lambda^2 + 10}}$$


$$\frac{1}{5} = \left| \frac{3\lambda - 8}{\sqrt{35} \cdot \sqrt{\lambda^2 + 10}} \right|$$

$$\text{Square } \Rightarrow \frac{1}{25} = \frac{9\lambda^2 + 64 - 48\lambda}{35(\lambda^2 + 10)}$$

$$\Rightarrow 19\lambda^2 - 120\lambda + 125 = 0$$

$$\Rightarrow 19\lambda^2 - 95\lambda - 25\lambda + 125 = 0$$

$$\Rightarrow x = 5, \frac{25}{19}$$



Perpendicular distance of point

$$(38\lambda_1, 10\lambda_2, 2) \equiv (50, 50, 2) \text{ from plane } P_1$$

$$= \frac{|3 \times 50 - 5 \times 50 + 2 - 7|}{\sqrt{35}} = \frac{105}{\sqrt{35}}$$

$$\text{Square } = \frac{105 \times 105}{35} = 315$$

84. Let $z = 1+i$ and $z_1 = \frac{1+i\bar{z}}{\bar{z}(1-z) + \frac{1}{z}}$. Then $\frac{12}{\pi} \arg(z_1)$ is equal to ____.

Official Ans. by NTA (9)

Allen Ans. (9)

$$\text{Sol. } z = 1+i$$

$$z_1 = \frac{1+i\bar{z}}{\bar{z}(1-z) + \frac{1}{z}}$$

$$z_1 = \frac{1+i(1-i)}{(1-i)(1-1-i) + \frac{1}{1+i}}$$

$$= \frac{1+i-i^2}{(1-i)(-i) + \frac{1-i}{2}}$$

$$= \frac{2+i}{-3i-1} = \frac{4+2i}{-3i-1}$$

$$= \frac{-(4+2i)(3i-1)}{(3i)^2 - (1)^2}$$

$$\text{Arg}(z_1) = \frac{3\pi}{4}$$

$$\therefore \frac{12}{\pi} \arg(z_1) = \frac{12}{\pi} \times \frac{3\pi}{4} = 9$$

85. $\lim_{x \rightarrow 0} \frac{48 \int_0^x \frac{t^3}{t^6 + 1} dt}{x^4}$ is equal to _____.

Official Ans. by NTA (12)

Allen Ans. (12)

Sol. $48 \lim_{x \rightarrow 0} \frac{\int_0^x \frac{t^3}{t^6 + 1} dt}{x^4} \left(\frac{0}{0} \right)$

Applying L' Hospitals Rule

$$48 \lim_{x \rightarrow 0} \frac{x^3}{x^6 + 1} \times \frac{1}{4x^3}$$

$$= 12$$

86. The mean and variance of 7 observations are 8 and 16 respectively. If one observation 14 is omitted a and b are respectively mean and variance of remaining 6 observation, then $a + 3b - 5$ is equal to _____.

Official Ans. by NTA (37)

Allen Ans. (37)

Sol. $\frac{x_1 + x_2 + \dots + x_7}{7} = 8$

$$\frac{x_1 + x_2 + x_3 + \dots + x_6 + 14}{7} = 8$$

$$\Rightarrow x_1 + x_2 + \dots + x_6 = 42$$

$$\therefore \frac{x_1 + x_2 + \dots + x_6}{6} = \frac{42}{6} = 7 = a$$

$$\frac{\sum x_i^2}{7} - 8^2 = 16$$

$$\Sigma x_i^2 = 560$$

$$\Rightarrow x_1^2 + x_2^2 + \dots + x_6^2 = 364$$

$$b = \frac{x_1^2 + x_2^2 + \dots + x_6^2}{6} - 7^2$$

$$= \frac{364}{6} - 49$$

$$b = \frac{70}{6}$$

$$a + 3b - 5 = 7 + 3 \times \frac{70}{6} - 5$$

$$= 37$$

87. If the equation of the plane passing through the point $(1, 1, 2)$ and perpendicular to the line $x - 3y + 2z - 1 = 0$ $4x - y + z$ is $Ax + By + Cz = 1$, then $140(C - B + A)$ is equal to _____.

Official Ans. by NTA (15)

Allen Ans. (15)

Sol. $x - 3y + 2z - 1 = 0$

$$4x - y + z = 0$$

$$\vec{n}_1 \times \vec{n}_2 = \begin{vmatrix} \hat{i} & \hat{j} & \hat{k} \\ 1 & -3 & 2 \\ 4 & -1 & 1 \end{vmatrix}$$

$$= -\hat{i} + 7\hat{j} + 11\hat{k}$$

Dr^s of normal to the plane is $-1, 7, 11$

Equation of plane :

$$-1(x - 1) + 7(y - 1) + 11(z - 2) = 0$$

$$-x + 7y + 11z = 28$$

$$\frac{-1}{28}x + \frac{7}{28}y + \frac{11}{28}z = 1$$

$$Ax + By + Cz = 1$$

$$140(C - B + A) = 140 \left(\frac{11}{28} - \frac{7}{28} - \frac{1}{28} \right)$$

$$= 140 \times \frac{3}{28} = 15$$

88. Let $\sum_{n=0}^{\infty} \frac{n^3((2n)!) + (2n-1)(n)!}{(n!)((2n)!)}$ = ae + $\frac{b}{e} + c$,

where $a, b, c \in \mathbb{Z}$ and $e = \sum_{n=0}^{\infty} \frac{1}{n!}$. Then $a^2 - b + c$ is

equal to _____.

Official Ans. by NTA (26)

Allen Ans. (26)

Sol.
$$\sum_{n=0}^{\infty} \frac{n^3((2n)!) + (2n-1)(n)!}{(n!)((2n)!)}$$

$$= \sum_{n=0}^{\infty} \frac{1}{(n-3)!} + \sum_{n=0}^{\infty} \frac{3}{(n-2)!}$$

$$+ \sum_{n=0}^{\infty} \frac{1}{(n-1)!} + \sum_{n=0}^{\infty} \frac{1}{(2n-1)!} - \sum_{n=0}^{\infty} \frac{1}{(2n)!}$$

$$= e + 3e + e + \frac{1}{2} \left(e - \frac{1}{e} \right) - \frac{1}{2} \left(e + \frac{1}{e} \right)$$

$$= 5e - \frac{1}{e}$$

$$\therefore a^2 - b + c = 26$$

- 89.** Number of 4-digit numbers (the repetition of digits is allowed) which are made using the digits 1, 2, 3 and 5, and are divisible by 15, is equal to _____

Official Ans. by NTA (21)

Allen Ans. (21)

Sol. For number to be divisible by 15, last digit should be 5 and sum of digits must be divisible by 3.

Possible combinations are

1	2	1	5
---	---	---	---

Numbers = 3

2	2	3	5
---	---	---	---

Numbers = 3

3	3	1	5
---	---	---	---

Numbers = 3

1	1	5	5
---	---	---	---

Numbers = 3

2	3	5	5
---	---	---	---

Numbers = 6

3	5	5	5
---	---	---	---

Numbers = 3

Total Numbers = 21

90. Let $f^1(x) = \frac{3x+2}{2x+3}, x \in \mathbb{R} - \left\{ \frac{-3}{2} \right\}$

For $n \geq 2$, define $f^n(x) = f^{10}f^{n-1}(x)$.

If $f^5(x) = \frac{ax+b}{bx+a}$, $\gcd(a,b)=1$, then $a+b$ is

equal to _____.

Official Ans. by NTA (3125)

Allen Ans. (3125)

Sol. $f^1(x) = \frac{3x+2}{2x+3}$

$$\Rightarrow f^2(x) = \frac{13x+12}{12x+13}$$

$$\Rightarrow f^3(x) = \frac{63x+62}{62x+63}$$

$$\therefore f^5(x) = \frac{1563x+1562}{1562x+1563}$$

$$a+b = 3125$$