

# FINAL JEE(Advanced) EXAMINATION - 2022

(Held On Sunday 28<sup>th</sup> AUGUST, 2022)

**PAPER-2**

**TEST PAPER WITH SOLUTION**

## CHEMISTRY

### SECTION-1 : (Maximum Marks : 24)

- This section contains **EIGHT (08)** questions.
- The answer to each question is a **SINGLE DIGIT INTEGER ranging from 0 TO 9, BOTH INCLUSIVE**.
- For each question, enter the correct integer corresponding to the answer using the mouse and the on-screen virtual numeric keypad in the place designated to enter the answer.
- Answer to each question will be evaluated according to the following marking scheme:


*Full Marks* : +3 If **ONLY** the correct integer is entered;

*Zero Marks* : 0 If the question is unanswered;

*Negative Marks* : -1 In all other cases.

1. Concentration of  $\text{H}_2\text{SO}_4$  and  $\text{Na}_2\text{SO}_4$  in a solution is 1 M and  $1.8 \times 10^{-2}$  M, respectively. Molar solubility of  $\text{PbSO}_4$  in the same solution is  $X \times 10^{-Y}$  M (expressed in scientific notation). The value of Y is \_\_\_\_\_.
- [Given: Solubility product of  $\text{PbSO}_4$  ( $K_{sp}$ ) =  $1.6 \times 10^{-8}$ . For  $\text{H}_2\text{SO}_4$ ,  $K_{a1}$  is very large and  $K_{a2} = 1.2 \times 10^{-2}$ ]


**Ans. (6)**



Since  $x$  is very small  $(1 + x) \approx 1$  and  $(1 - x) \approx 1$

$$x = (1.8 \times 10^{-2} - 1.2 \times 10^{-2}) M$$

$$\begin{aligned} [\text{SO}_4^{2-}] &= (1.8 \times 10^{-2} - 0.6 \times 10^{-2}) M \\ &= 1.2 \times 10^{-2} M \end{aligned}$$



$$K_{sp} = s(s + 1.2 \times 10^{-2}) = 1.6 \times 10^{-8}$$

(PbSO<sub>4</sub>)

Here,  $(s + 1.2 \times 10^{-2}) \approx 1.2 \times 10^{-2}$  (since 's' is very small)

$$s(1.2 \times 10^{-2}) = 1.6 \times 10^{-8}$$

$$\Rightarrow s = \frac{1.6}{1.2} \times 10^{-6} M = X \times 10^{-Y} M$$

$$\Rightarrow Y = 6$$

2. An aqueous solution is prepared by dissolving 0.1 mol of an ionic salt in 1.8 kg of water at 35 °C. The salt remains 90% dissociated in the solution. The vapour pressure of the solution is 59.724 mm of Hg. Vapor pressure of water at 35 °C is 60.000 mm of Hg. The number of ions present per formula unit of the ionic salt is \_\_\_\_\_.


**Ans. (5)**

**Sol.** 0.1 mole ionic salt in 1.8 kg water at 35° C

Vapour pressure of solution = 59.724 mm of Hg

Vapour pressure of pure H<sub>2</sub>O = 60.000 mm of Hg

Let the number of ions present per formula unit of the ionic salt be 'x'



Total moles of non-volatile particles =  $0.01 + 0.09x$   
in 1.8 kg water

$$\text{Moles of water} = \frac{1.8 \times 10^3}{18} = 100 \text{ moles}$$

$$\text{Relative lowering of vapour pressure} \frac{P^\circ - P_s}{P^\circ} = \text{Mole fraction of non - volatile particles}$$

$$\frac{P^\circ - P_s}{P_s} = \frac{\text{moles of non-volatile particles}}{\text{moles of water}}$$

$$\frac{60.000 - 59.724}{59.724} = \frac{0.01 + 0.09x}{100}$$

$$(0.276) \times 100 = 0.59274 + (0.59274 \times 9)x$$

$$27.6 - 0.59274 = (0.59274 \times 9)x$$

$$\Rightarrow x \approx \frac{27}{0.6 \times 9} = 5$$


3. Consider the strong electrolytes  $Z_mX_n$ ,  $U_mY_p$  and  $V_mX_n$ . Limiting molar conductivity ( $\Lambda^\circ$ ) of  $U_mY_p$  and  $V_mX_n$  are 250 and 440  $S\text{ cm}^2\text{ mol}^{-1}$ , respectively. The value of  $(m + n + p)$  is \_\_\_\_\_.

Given:

Ion	$Z^{n+}$	$U^{p+}$	$V^{n+}$	$X^{m-}$	$Y^{m-}$
$\lambda^\circ (S\text{ cm}^2\text{ mol}^{-1})$	50.0	25.0	100.0	80.0	100.0

$\lambda^\circ$  is the limiting molar conductivity of ions

The plot of molar conductivity ( $\Lambda$ ) of  $Z_mX_n$  vs  $c^{1/2}$  is given below.



Ans. (7)

Sol.  $\Lambda^\circ(U_mY_p) = m \times \lambda^\circ_{U^{p+}} + p \times \lambda^\circ_{Y^{m-}} = 250$


$$25m + 100p = 250$$

$$m + 4p = 10 \quad \dots\dots(1)$$

$$\Lambda^\circ(V_mX_n) = m \times \lambda^\circ_{V^{n+}} + n \times \lambda^\circ_{X^{m-}} = 440$$

$$100m + 80n = 440$$

$$5m + 4n = 22 \quad \dots\dots(2)$$



From the extrapolation of curve

$$\Delta^{\circ}(Z_m X_n) = 340$$

$$m \times \lambda_{Z^{m+}}^{\circ} + n \lambda_{X^{n-}}^{\circ} = 340$$

$$50m + 80n = 340$$

$$5m + 8n = 34 \quad \dots\dots(3)$$

$$(3) - (2) \Rightarrow 4n = 12 \Rightarrow n = 3$$



Putting in (2) we get  $m = 2$

Putting in (1) we get  $p = 2$

$$m + n + p = 2 + 3 + 2 = 7$$

4. The reaction of Xe and  $O_2F_2$  gives a Xe compound **P**. The number of moles of HF produced by the complete hydrolysis of 1 mol of **P** is \_\_\_\_\_.

**Ans. (4)**



$\therefore$  One mole of  $XeF_4$  gives 4 moles of HF on hydrolysis.

5. Thermal decomposition of  $AgNO_3$  produces two paramagnetic gases. The total number of electrons present in the antibonding molecular orbitals of the gas that has the higher number of unpaired electrons is \_\_\_\_\_.


**Ans. (6)**



– Both  $NO_2$  &  $O_2$  are paramagnetic


–  $NO_2$  is odd electron molecule with one unpaired electron

–  $O_2$  has two unpaired electrons




Total number of antibonding electrons = 6


6. The number of isomeric tetraenes (NOT containing  $sp$ -hybridized carbon atoms) that can be formed from the following reaction sequence is \_\_\_\_\_.




Ans. (2)




7. The number of  $-CH_2-$  (methylene) groups in the product formed from the following reaction sequence is \_\_\_\_\_.



Ans. (0)




8. The total number of chiral molecules formed from one molecule of **P** on complete ozonolysis ( $O_3, Zn/H_2O$ ) is \_\_\_\_\_.



Ans. (2)

Sol.



SECTION-2 : (Maximum Marks : 24)

- This section contains **SIX (06)** questions.
- Each question has **FOUR** options (A), (B), (C) and (D). **ONE OR MORE THAN ONE** of these four option(s) is (are) correct answer(s).
- For each question, choose the option(s) corresponding to (all) the correct answer(s).
- Answer to each question will be evaluated according to the following marking scheme:

*Full Marks* : +4 **ONLY** if (all) the correct option(s) is(are) chosen;

*Partial Marks* : +3 If all the four options are correct but **ONLY** three options are chosen;

*Partial Marks* : +2 If three or more options are correct but **ONLY** two options are chosen, both of which are correct;

*Partial Marks* : +1 If two or more options are correct but **ONLY** one option is chosen and it is a correct option;

*Zero Marks* : 0 If unanswered;

*Negative Marks* : -2 In all other cases.

9. To check the principle of multiple proportions, a series of pure binary compounds ( $P_mQ_n$ ) were analyzed and their composition is tabulated below. The correct option(s) is(are)

Compound	Weight % of P	Weight % of Q
1	50	50
2	44.4	55.6
3	40	60

- (A) If empirical formula of compound 3 is  $P_3Q_4$ , then the empirical formula of compound 2 is  $P_3Q_5$ .  
 (B) If empirical formula of compound 3 is  $P_3Q_2$  and atomic weight of element P is 20, then the atomic weight of Q is 45.  
 (C) If empirical formula of compound 2 is PQ, then the empirical formula of the compound 1 is  $P_5Q_4$ .  
 (D) If atomic weight of P and Q are 70 and 35, respectively, then the empirical formula of compound 1 is  $P_2Q$ .

Ans. (B,C)

Sol.

Compound	Weight % of P	Weight % of Q
1	50	50
2	44.4	55.6
3	40	60

For option (A)

Let atomic mass of P be  $M_P$  and atomic mass of Q be  $M_Q$

Molar ratio of atoms P : Q in compound 3 is

$$\frac{40}{M_P} : \frac{60}{M_Q} = 3 : 4$$

$$\frac{2M_Q}{3M_P} = \frac{3}{4} \Rightarrow 9M_P = 8M_Q$$

Molar ratio of atoms P : Q in compound 2 is

$$\begin{aligned} & \frac{44.4}{M_P} : \frac{55.6}{M_Q} \\ &= 44.4 M_Q : 55.6 M_P \\ &= 44.4 M_Q : 55.6 \times \frac{8M_Q}{9} \\ &= 44.4 : 55.6 \times \frac{8}{9} \\ &= 9 : 10 \end{aligned}$$

$\Rightarrow$  Empirical formula of compound 2 is therefore  $P_9Q_{10}$

Option (A) is incorrect

For option (B)



Molar Ratio of atoms P : Q in compound 3 is  $\frac{40}{M_p} : \frac{60}{M_Q} = 3 : 2$

$$\frac{2M_Q}{3M_p} = \frac{3}{2} \Rightarrow 9M_p = 4M_Q$$

If  $M_p = 20 \Rightarrow M_Q = \frac{9 \times 20}{4} = 45$

Option (B) is correct

For option (C)

Molar ratio of atoms P : Q in compound 2 is

$$\frac{44.4}{M_p} : \frac{55.6}{M_Q} = 44.4M_Q : 55.6M_p = 1 : 1$$

$$\Rightarrow \frac{M_p}{M_Q} = \frac{44.4}{55.6}$$

Molar ratio of atoms P : Q in compound 1 is

$$\frac{50}{M_p} : \frac{50}{M_Q} = M_Q : M_p$$

$$= 55.6 : 44.4$$

$$\approx 5 : 4$$

Hence, empirical formula of compound 1 is  $P_5Q_4$

Hence, option (C) is correct

For option (D)

Molar ratio of atoms P : Q in compound 1 is

$$\frac{50}{M_p} : \frac{50}{M_Q} = M_Q : M_p$$

$$= 35 : 70 = 1 : 2$$

Hence, empirical formula of compound 1 is  $PQ_2$

Hence, option (D) is incorrect

10. The correct option(s) about entropy (S) is(are)

[R = gas constant, F = Faraday constant, T = Temperature]

(A) For the reaction,  $M(s) + 2H^+(aq) \rightarrow H_2(g) + M^{2+}(aq)$ , if  $\frac{dE_{\text{cell}}}{dT} = \frac{R}{F}$ , then the entropy change of the reaction is R (assume that entropy and internal energy changes are temperature independent).

(B) The cell reaction,  $Pt(s) | H_2(g, 1\text{bar}) | H^+(aq, 0.01M) || H^+(aq, 0.1M) | H_2(g, 1\text{bar}) | Pt(s)$ , is an entropy driven process.

(C) For racemization of an optically active compound,  $\Delta S > 0$ .

(D)  $\Delta S > 0$ , for  $[Ni(H_2O)_6]^{2+} + 3\text{en} \rightarrow [Ni(\text{en})_3]^{2+} + 6H_2O$  (where en = ethylenediamine).

Ans. (B,C,D)

Sol.  $\Delta G = \Delta H - T\Delta S$

$$\Delta G = \Delta H + T \left( \frac{d\Delta G}{dT} \right)_p$$

$$-nF \left( \frac{dE_{\text{cell}}}{dT} \right) = -\Delta S$$

$$\frac{dE_{\text{cell}}}{dT} = \frac{\Delta S}{nF} = \frac{R}{F} \text{ (given)}$$


$$\Rightarrow \Delta S = nR$$



$$n = 2$$

$$\Rightarrow \Delta S = 2R$$

Hence, option (A) is incorrect



$$E_{\text{cell}} = E^{\circ}_{\text{cell}} - \frac{0.0591}{1} \log \frac{0.01}{0.1} = 0.0591V$$

$E_{\text{cell}}$  is positive  $\Rightarrow \Delta G < 0$  and  $\Delta S > 0$  ( $\Delta H = 0$  for concentration cells)


Hence, option (B) is correct

Racemization of an optically active compound is a spontaneous process.

Here,  $\Delta H = 0$  (similar type of bonds are present in enantiomers)

$$\Rightarrow \Delta S > 0$$

Hence, option (C) is correct.



more stable complex is formed

$$\Rightarrow \Delta S > 0$$

Hence, option (D) is correct.




11. The compound(s) which react(s) with  $NH_3$  to give boron nitride (BN) is(are)

- (A) B                      (B)  $B_2H_6$                       (C)  $B_2O_3$                       (D)  $BF_4$

Ans. (B,C)



Boron produced BN with ammonia but **Boron is element not compound**. So that this option not involve in answer.




12. The correct option(s) related to the extraction of iron from its ore in the blast furnace operating in the temperature range 900 – 1500 K is(are)
- (A) Limestone is used to remove silicate impurity.
- (B) Pig iron obtained from blast furnace contains about 4% carbon.
- (C) Coke (C) converts  $\text{CO}_2$  to  $\text{CO}$ .
- (D) Exhaust gases consist of  $\text{NO}_2$  and  $\text{CO}$ .

Ans. (A,B,C)


- Sol. (A)  $\text{CaO} + \text{SiO}_2 \rightarrow \text{CaSiO}_3$  (in the temperature range 900 – 1500 K)
- (B) In fusion zone molten iron becomes heavy by absorbing elemental impurities and produces Pig iron. (in the temperature range 900 – 1500 K)
- (C)  $\text{C} + \text{CO}_2 \rightarrow 2\text{CO}$  (in the temperature range 900 – 1500 K)
- (D) Exhaust gases does not contain  $\text{NO}_2$ .

13. Considering the following reaction sequence, the correct statement(s) is(are)



- (A) Compounds **P** and **Q** are carboxylic acids.
- (B) Compound **S** decolorizes bromine water.
- (C) Compounds **P** and **S** react with hydroxylamine to give the corresponding oximes.
- (D) Compound **R** reacts with dialkylcadmium to give the corresponding tertiary alcohol.

Ans. (A,C)



14. Among the following, the correct statement(s) about polymers is(are)
- (A) The polymerization of chloroprene gives natural rubber.
- (B) Teflon is prepared from tetrafluoroethene by heating it with persulphate catalyst at high pressures.
- (C) PVC are thermoplastic polymers.
- (D) Ethene at 350-570 K temperature and 1000-2000 atm pressure in the presence of a peroxide initiator yields high density polythene.

Ans. (B,C)

Sol. (a) The polymerisation of neoprene gives natural rubber.

(b) is correct statement

(c) is correct statement

(d) Ethene at 350-570 K temperature and 1000-2000 atm pressure in the presence of a peroxide initiator yields low density polythene.

**SECTION-3 : (Maximum Marks : 12)**

- This section contains **FOUR (04)** questions.
- Each question has **FOUR** options (A), (B), (C) and (D). **ONLY ONE** of these four options is the correct answer.
- For each question, choose the option corresponding to the correct answer.
- Answer to each question will be evaluated according to the following marking scheme:  
*Full Marks* : +3 If **ONLY** the correct option is chosen;  
*Zero Marks* : 0 If none of the options is chosen (i.e. the question is unanswered);  
*Negative Marks* : -1 In all other cases.

- 15.** Atom X occupies the fcc lattice sites as well as alternate tetrahedral voids of the same lattice. The packing efficiency (in %) of the resultant solid is closest to  
 (A) 25                      (B) 35                      (C) 55                      (D) 75

**Ans. (B)**

Atom 'X' occupies FCC lattice points as well as alternate tetrahedral voids of the same lattice

$\Rightarrow \frac{1}{4}$ th distance of body diagonal

$$= \frac{\sqrt{3}a}{4} = 2r_x$$

$$\Rightarrow a = \frac{8r_x}{\sqrt{3}}$$

Number of atoms of X per unit cell

$$= 4 + 4 = 8$$

(FCC lattice points)

(Alternate tetrahedral voids)

$$\% \text{ packing efficiency} = \frac{\text{Volume occupied by X}}{\text{Volume of cubic unit cell}} \times 100$$

$$= \frac{8 \times \frac{4}{3} \pi (r_x)^3}{a^3} \times 100$$

$$= \frac{8 \times \frac{4}{3} \pi (r_x)^3}{\left(\frac{8r_x}{\sqrt{3}}\right)^3} \times 100$$

$$= \left(8 \times \frac{4}{3} \times \pi \times \frac{1}{8^3} \times 3\sqrt{3}\right) \times 100$$



$$= \frac{\sqrt{3}\pi}{16} \times 100$$

$$= 34\%$$

Hence, option (B) is the most appropriate option

16. The reaction of  $\text{HClO}_3$  with  $\text{HCl}$  gives a paramagnetic gas, which upon reaction with  $\text{O}_3$  produces  
 (A)  $\text{Cl}_2\text{O}$  (B)  $\text{ClO}_2$  (C)  $\text{Cl}_2\text{O}_6$  (D)  $\text{Cl}_2\text{O}_7$



Ans. (C)





17. The reaction  $\text{Pb}(\text{NO}_3)_2$  and  $\text{NaCl}$  in water produces a precipitate that dissolves upon the addition of  $\text{HCl}$  of appropriate concentration. The dissolution of the precipitate is due to the formation of



- (A)  $\text{PbCl}_2$  (B)  $\text{PbCl}_4$  (C)  $[\text{PbCl}_4]^{2-}$  (D)  $[\text{PbCl}_6]^{2-}$

Ans. (C)



18. Treatment of D-glucose with aqueous  $\text{NaOH}$  results in a mixture of monosaccharides, which are





Ans. (C)

Sol. Basic catalyse tautomerism through enediol intermediate

