





**FINAL JEE(Advanced) EXAMINATION – 2023****(Held On Sunday 04<sup>th</sup> June, 2023)****PAPER-1****TEST PAPER****CHEMISTRY****SECTION-1 : (Maximum Marks : 12)**

- This section contains **THREE (03)** questions.
- Each question has **FOUR** options (A), (B), (C) and (D). **ONE OR MORE THAN ONE** of these four option(s) is(are) correct answer(s).
- For each question, choose the option(s) corresponding to (all) the correct answer(s).
- Answer to each question will be evaluated according to the following marking scheme:
  - Full Marks* : +4 **ONLY** if (all) the correct option(s) is(are) chosen;
  - Partial Marks* : +3 If all the four options are correct but **ONLY** three options are chosen;
  - Partial Marks* : +2 If three or more options are correct but **ONLY** two options are chosen, both of which are correct;
  - Partial Marks* : +1 If two or more options are correct but **ONLY** one option is chosen and it is a correct option;
  - Zero Marks* : 0 If none of the options is chosen (i.e. the question is unanswered);
  - Negative Marks* : -2 In all other cases.
- For example, in a question, if (A), (B) and (D) are the **ONLY** three options corresponding to correct answers, then
  - choosing **ONLY** (A), (B) and (D) will get +4 marks;
  - choosing **ONLY** (A) and (B) will get +2 marks;
  - choosing **ONLY** (A) and (D) will get +2 marks;
  - choosing **ONLY** (B) and (D) will get +2 marks;
  - choosing **ONLY** (A) will get +1 marks;
  - choosing **ONLY** (B) will get +1 marks;
  - choosing **ONLY** (D) will get +1 marks;
  - choosing no option (i.e. the question is unanswered) will get 0 marks; and
  - choosing any other combination of options will get -2 marks.

1. The correct statement(s) related to processes involved in the extraction of metals is(are)
  - (A) Roasting of Malachite produces Cuprite.
  - (B) Calcination of Calamine produces Zincite.
  - (C) Copper pyrites is heated with silica in a reverberatory furnace to remove iron.
  - (D) Impure silver is treated with aqueous KCN in the presence of oxygen followed by reduction with zinc metal.



2. In the following reactions, **P**, **Q**, **R**, and **S** are the major products.















The correct statement(s) about **P**, **Q**, **R**, and **S** is(are)

- (A) Both **P** and **Q** have asymmetric carbon(s).  
 (B) Both **Q** and **R** have asymmetric carbon(s).  
 (C) Both **P** and **R** have asymmetric carbon(s).  
 (D) **P** has asymmetric carbon(s), **S** does **not** have any asymmetric carbon.

3. Consider the following reaction scheme and choose the correct option(s) for the major products **Q**, **R** and **S**.




(A)    (B)   

(C)    (D)   

SECTION-2 : (Maximum Marks : 12)

- This section contains **FOUR (04)** questions.
- Each question has **FOUR** options (A), (B), (C) and (D). **ONLY ONE** of these four options is the correct answer.
- For each question, choose the option corresponding to the correct answer.
- Answer to each question will be evaluated according to the following marking scheme:  
*Full Marks* : +3 If **ONLY** the correct option is chosen;  
*Zero Marks* : 0 If none of the options is chosen (i.e. the question is unanswered);  
*Negative Marks* : -1 In all other cases.

4. In the scheme given below, **X** and **Y**, respectively, are




- (A)  $\text{CrO}_4^{2-}$  and  $\text{Br}_2$  (B)  $\text{MnO}_4^{2-}$  and  $\text{Cl}_2$   
 (C)  $\text{MnO}_4^-$  and  $\text{Cl}_2$  (D)  $\text{MnSO}_4$  and  $\text{HOCl}$
5. Plotting  $1/\Lambda_m$  against  $c\Lambda_m$  for aqueous solutions of a monobasic weak acid (HX) resulted in a straight line with y-axis intercept of P and slope of S. The ratio P/S is  
 [ $\Lambda_m$  = molar conductivity  
 $\Lambda_m^\circ$  = limiting molar conductivity  
 $c$  = molar concentration  
 $K_a$  = dissociation constant of HX]  
 (A)  $K_a \Lambda_m^\circ$  (B)  $K_a \Lambda_m^\circ / 2$  (C)  $2 K_a \Lambda_m^\circ$  (D)  $1 / (K_a \Lambda_m^\circ)$
6. On decreasing the pH from 7 to 2, the solubility of a sparingly soluble salt (MX) of a weak acid (HX) increased from  $10^{-4} \text{ mol L}^{-1}$  to  $10^{-3} \text{ mol L}^{-1}$ . The  $pK_a$  of HX is:  
 (A) 3 (B) 4 (C) 5 (D) 2
7. In the given reaction scheme, **P** is a phenyl alkyl ether, **Q** is an aromatic compound; **R** and **S** are the major products.
- $$\text{P} \xrightarrow{\text{HI}} \text{Q} \xrightarrow[\text{(iii) H}_3\text{O}^+]{\text{(i) NaOH, (ii) CO}_2} \text{R} \xrightarrow[\text{(ii) H}_3\text{O}^+]{\text{(i) (CH}_3\text{CO)}_2\text{O}} \text{S}$$
- The correct statement about **S** is  
 (A) It primarily inhibits noradrenaline degrading enzymes.  
 (B) It inhibits the synthesis of prostaglandin.  
 (C) It is a narcotic drug.  
 (D) It is *ortho*-acetylbenzoic acid.

**SECTION-3 : (Maximum Marks : 24)**

- This section contains **SIX (06)** questions.
- The answer to each question is a **NON-NEGATIVE INTEGER**.
- For each question, enter the correct integer corresponding to the answer using the mouse and the on-screen virtual numeric keypad in the place designated to enter the answer.
- Answer to each question will be evaluated according to the following marking scheme:  

Full Marks	: +4	ONLY If the correct integer is entered;
Zero Marks	: 0	In all other cases.

8. The stoichiometric reaction of 516 g of dimethyldichlorosilane with water results in a tetrameric cyclic product **X** in 75% yield. The weight (in g) of **X** obtained is \_\_\_\_.  
 [Use, molar mass ( $\text{g mol}^{-1}$ ): H = 1, C = 12, O = 16, Si = 28, Cl = 35.5]
9. A gas has a compressibility factor of 0.5 and a molar volume of  $0.4 \text{ dm}^3 \text{ mol}^{-1}$  at a temperature of 800 K and pressure  $x \text{ atm}$ . If it shows ideal gas behaviour at the same temperature and pressure, the molar volume will be  $y \text{ dm}^3 \text{ mol}^{-1}$ . The value of  $x/y$  is \_\_\_\_.  
 [Use: Gas constant,  $R = 8 \times 10^{-2} \text{ L atm K}^{-1} \text{ mol}^{-1}$ ]
10. The plot of  $\log k_f$  versus  $1/T$  for a reversible reaction  $A(g) \rightleftharpoons P(g)$  is shown.




Pre-exponential factors for the forward and backward reactions are  $10^{15} \text{ s}^{-1}$  and  $10^{11} \text{ s}^{-1}$ , respectively. If the value of  $\log K$  for the reaction at 500 K is 6, the value of  $|\log k_b|$  at 250 K is \_\_\_\_.

[K = equilibrium constant of the reaction

$k_f$  = rate constant of forward reaction

$k_b$  = rate constant of backward reaction]


11. One mole of an ideal monoatomic gas undergoes two reversible processes ( $A \rightarrow B$  and  $B \rightarrow C$ ) as shown in the given figure :



$A \rightarrow B$  is an adiabatic process. If the total heat absorbed in the entire process ( $A \rightarrow B$  and  $B \rightarrow C$ ) is  $RT_2 \ln 10$ , the value of  $2 \log V_3$  is \_\_\_\_\_.

[Use, molar heat capacity of the gas at constant pressure,  $C_{p,m} = \frac{5}{2} R$ ]


12. In a one-litre flask, 6 moles of A undergoes the reaction  $A(g) \rightleftharpoons P(g)$ . The progress of product formation at two temperatures (in Kelvin),  $T_1$  and  $T_2$ , is shown in the figure:



If  $T_1 = 2T_2$  and  $(\Delta G_2^\ominus - \Delta G_1^\ominus) = RT_2 \ln x$ , then the value of  $x$  is \_\_\_\_\_.

[ $\Delta G_1^\ominus$  and  $\Delta G_2^\ominus$  are standard Gibb's free energy change for the reaction at temperatures  $T_1$  and  $T_2$ , respectively.]

13. The total number of  $sp^2$  hybridised carbon atoms in the major product **P** (a non-heterocyclic compound) of the following reaction is \_\_\_\_\_.



**SECTION-4 : (Maximum Marks : 12)**

- This section contains **FOUR (04)** Matching List Sets.
- Each set has **ONE** Multiple Choice Question.
- Each set has **TWO** lists : **List-I** and **List-II**.
- **List-I** has **Four** entries (P), (Q), (R) and (S) and **List-II** has **Five** entries (1), (2), (3), (4) and (5).
- **FOUR** options are given in each Multiple Choice Question based on **List-I** and **List-II** and **ONLY ONE** of these four options satisfies the condition asked in the Multiple Choice Question.
- Answer to each question will be evaluated according to the following marking scheme:  
*Full Marks* : +3 **ONLY** if the option corresponding to the correct combination is chosen;  
*Zero Marks* : 0 If none of the options is chosen (i.e. the question is unanswered);  
*Negative Marks* : -1 In all other cases.

14. Match the reactions (in the given stoichiometry of the reactants) in List-I with one of their products given in List-II and choose the correct option.

**List-I**

- (P)  $\text{P}_2\text{O}_3 + 3\text{H}_2\text{O} \rightarrow$   
 (Q)  $\text{P}_4 + 3\text{NaOH} + 3\text{H}_2\text{O} \rightarrow$   
 (R)  $\text{PCl}_5 + \text{CH}_3\text{COOH} \rightarrow$   
 (S)  $\text{H}_3\text{PO}_2 + 2\text{H}_2\text{O} + 4\text{AgNO}_3 \rightarrow$

**List-II**

- (1)  $\text{P(O)(OCH}_3\text{)Cl}_2$   
 (2)  $\text{H}_3\text{PO}_3$   
 (3)  $\text{PH}_3$   
 (4)  $\text{POCl}_3$   
 (5)  $\text{H}_3\text{PO}_4$

- (A)  $\text{P} \rightarrow 2; \text{Q} \rightarrow 3; \text{R} \rightarrow 1; \text{S} \rightarrow 5$   
 (B)  $\text{P} \rightarrow 3; \text{Q} \rightarrow 5; \text{R} \rightarrow 4; \text{S} \rightarrow 2$   
 (C)  $\text{P} \rightarrow 5; \text{Q} \rightarrow 2; \text{R} \rightarrow 1; \text{S} \rightarrow 3$   
 (D)  $\text{P} \rightarrow 2; \text{Q} \rightarrow 3; \text{R} \rightarrow 4; \text{S} \rightarrow 5$

15. Match the electronic configurations in List-I with appropriate metal complex ions in List-II and choose the correct option.

[Atomic Number: Fe = 26, Mn = 25, Co = 27]

**List-I**


- (P)  $t_{2g}^6 e_g^0$   
 (Q)  $t_{2g}^3 e_g^2$   
 (R)  $e^2 t_2^3$   
 (S)  $t_{2g}^4 e_g^2$

**List-II**

- (1)  $[\text{Fe}(\text{H}_2\text{O})_6]^{2+}$   
 (2)  $[\text{Mn}(\text{H}_2\text{O})_6]^{2+}$   
 (3)  $[\text{Co}(\text{NH}_3)_6]^{3+}$   
 (4)  $[\text{FeCl}_4]^-$   
 (5)  $[\text{CoCl}_4]^{2-}$

- (A)  $\text{P} \rightarrow 1; \text{Q} \rightarrow 4; \text{R} \rightarrow 2; \text{S} \rightarrow 3$   
 (B)  $\text{P} \rightarrow 1; \text{Q} \rightarrow 2; \text{R} \rightarrow 4; \text{S} \rightarrow 5$   
 (C)  $\text{P} \rightarrow 3; \text{Q} \rightarrow 2; \text{R} \rightarrow 5; \text{S} \rightarrow 1$   
 (D)  $\text{P} \rightarrow 3; \text{Q} \rightarrow 2; \text{R} \rightarrow 4; \text{S} \rightarrow 1$

16. Match the reactions in List-I with the features of their products in List-II and choose the correct option.

- | List-I  | List-II                           |
|---|-----------------------------------|
| (P) (-)-1-Bromo-2-ethylpentane $\xrightarrow[\text{S}_\text{N}2 \text{ reaction}]{\text{aq. NaOH}}$<br>(single enantiomer)  | (1) Inversion of configuration    |
| (Q) (-)-2-Bromopentane $\xrightarrow[\text{S}_\text{N}2 \text{ reaction}]{\text{aq. NaOH}}$<br>(single enantiomer)  | (2) Retention of configuration    |
| (R) (-)-3-Bromo-3-methylhexane $\xrightarrow[\text{S}_\text{N}1 \text{ reaction}]{\text{aq. NaOH}}$<br>(single enantiomer)  | (3) Mixture of enantiomers        |
| (S) <br>(Single enantiomer) $\xrightarrow[\text{S}_\text{N}1 \text{ reaction}]{\text{aq. NaOH}}$ | (4) Mixture of structural isomers |

- (A) P  $\rightarrow$  1; Q  $\rightarrow$  2; R  $\rightarrow$  5; S  $\rightarrow$  3  
(B) P  $\rightarrow$  2; Q  $\rightarrow$  1; R  $\rightarrow$  3; S  $\rightarrow$  5  
(C) P  $\rightarrow$  1; Q  $\rightarrow$  2; R  $\rightarrow$  5; S  $\rightarrow$  4  
(D) P  $\rightarrow$  2; Q  $\rightarrow$  4; R  $\rightarrow$  3; S  $\rightarrow$  5

17. The major products obtained from the reactions in List-II are the reactants for the named reactions mentioned in List-I. Match List-I with List-II and choose the correct option.

- | List-I                       | List-II   |
|------------------------------|---|
| (P) Etard reaction           | (1) Acetophenone $\xrightarrow{\text{Zn-Hg, HCl}}$                                    |
| (Q) Gattermann reaction      | (2) Toluene $\xrightarrow[\text{(ii) SOCl}_2]{\text{(i) KMnO}_4, \text{KOH}, \Delta}$ |
| (R) Gattermann-Koch reaction | (3) Benzene $\xrightarrow[\text{anhyd. AlCl}_3]{\text{CH}_3\text{Cl}}$                |
| (S) Rosenmund reduction      | (4) Aniline $\xrightarrow[273-278 \text{ K}]{\text{NaNO}_2/\text{HCl}}$               |
|                              | (5) Phenol $\xrightarrow{\text{Zn}, \Delta}$  |

- (A) P  $\rightarrow$  2; Q  $\rightarrow$  4; R  $\rightarrow$  1; S  $\rightarrow$  3  
(B) P  $\rightarrow$  1; Q  $\rightarrow$  3; R  $\rightarrow$  5; S  $\rightarrow$  2  
(C) P  $\rightarrow$  3; Q  $\rightarrow$  2; R  $\rightarrow$  1; S  $\rightarrow$  4  
(D) P  $\rightarrow$  3; Q  $\rightarrow$  4; R  $\rightarrow$  5; S  $\rightarrow$  2