2.


JEE Advanced Chemistry 10 Years Topicwise Questions with Solutions

ORGANIC CHEMISTRY

ALCOHOL, PHENOL & ETHER

1. In the following reaction sequence, the correct structure(s) of X is (are)

[JEE(Advanced) 2018]



[JEE(Advanced) 2016]

Cumene(C₉H₁₂) $\xrightarrow{(i) O_2}$ P $\xrightarrow{CHCl_3/NaOH}$ Q (major) + R (minor)

$$Q \xrightarrow{\text{NaOH}} S$$


- (A) **R** is steam volatile
- (B) \mathbf{Q} gives dark violet coloration with 1% aqueous FeCl₃ solution
- (C) S gives yellow precipitate with 2, 4,-dinitrophenylhydrazine
- (D) S gives dark violet coloration with 1% aqueous $FeCl_3$ solution
- **3.** The number of hydroxyl group(s) in **Q** is

 $\begin{array}{c} H \\ HO \\ HO \\ H_{3}C \\ CH_{3} \end{array} \xrightarrow{H^{+}} \mathbf{P} \xrightarrow{\text{aqueous dilute KMNO}_{4}(\text{excess})} \mathbf{Q} \\ \end{array} \xrightarrow{0^{\circ}C} \mathbf{Q}$

[JEE(Advanced) 2015]



4. The acidic hydrolysis of ether (X) shown below is fastest when




(A) one phenyl group is replaced by a methyl group

- (B) one phenyl group is replaced by a para-methoxyphenyl group
- (C) two phenyl groups are replaced by two para-methoxyphenyl group
- (D) no structural change is made to \boldsymbol{X}

1


SOLUTIONS




Q gives dark violet coloration with 1% aqueous FeCl₃ solution because it has phenolic (–OH) group.



3.



4. Ans. (C)



- groups then carbocation formed in above sequence is If 2 Ph groups are substituted by 2 MeO- $\langle O \rangle$ more stable and rate of above hydrolysis increases.