

PHYSICAL CHEMISTRY

MOLE CONCEPT

1. The treatment of an aqueous solution of 3.74 g of $\text{Cu}(\text{NO}_3)_2$ with excess KI results in a brown solution along with the formation of a precipitate. Passing H_2S through this brown solution gives another precipitate X. The amount of X (in g) is _____.

[Given : Atomic mass of H = 1, N = 14, O = 16, S = 32, K = 39, Cu = 63, I = 127]

[JEE(Advanced) 2022]

2. Dissolving 1.24 g of white phosphorous in boiling NaOH solution in an inert atmosphere gives a gas Q. The amount of CuSO_4 (in g) required to completely consume the gas Q is _____.

[Given : Atomic mass of H = 1, O = 16, Na = 23, P = 31, S = 32, Cu = 63]

[JEE(Advanced) 2022]

3. To check the principle of multiple proportions, a series of pure binary compounds (P_mQ_n) were analyzed and their composition is tabulated below. The correct option(s) is(are)

[JEE(Advanced) 2022]

Compound	Weight % of P	Weight % of Q
1	50	50
2	44.4	55.6
3	40	60

(A) If empirical formula of compound 3 is P_3Q_4 , then the empirical formula of compound 2 is P_3Q_5 .

(B) If empirical formula of compound 3 is P_3Q_2 and atomic weight of element P is 20, then the atomic weight of Q is 45.

(C) If empirical formula of compound 2 is PQ, then the empirical formula of the compound 1 is P_5Q_4 .

(D) If atomic weight of P and Q are 70 and 35, respectively, then the empirical formula of compound 1 is P_2Q .

Question Stem for Question Nos. 4 and 5

Question Stem

Reaction of x g of Sn with HCl quantitatively produced a salt. Entire amount of the salt reacted with y g of nitrobenzene in the presence of required amount of HCl to produce 1.29 g of an organic salt (quantitatively).

(Use Molar masses (in g mol^{-1}) of H, C, N, O, Cl and Sn as 1, 12, 14, 16, 35 and 119, respectively).

[JEE(Advanced) 2021]

4. The value of x is _____.
5. The value of y is _____.

6. Aluminium reacts with sulfuric acid to form aluminium sulfate and hydrogen. What is the volume of hydrogen gas in liters (L) produced at 300 K and 1.0 atm pressure, when 5.4 g of aluminium and 50.0 mL of 5.0 M sulfuric acid are combined for the reaction ?

(Use molar mass of aluminium as 27.0 g mol^{-1} , $R = 0.082 \text{ atm L mol}^{-1} \text{ K}^{-1}$) [JEE(Advanced) 2020]

7. The ammonia prepared by treating ammonium sulphate with calcium hydroxide is completely used by $\text{NiCl}_2 \cdot 6\text{H}_2\text{O}$ to form a stable coordination compound. Assume that both the reactions are 100% complete. If 1584 g of ammonium sulphate and 952g of $\text{NiCl}_2 \cdot 6\text{H}_2\text{O}$ are used in the preparation, the combined weight (in grams) of gypsum and the nickel-ammonia coordination compound thus produced is _____.

(Atomic weights in g mol^{-1} : H = 1, N = 14, O = 16, S = 32, Cl = 35.5, Ca = 40, Ni = 59)


[JEE(Advanced) 2018]

8. If the value of Avogadro number is $6.023 \times 10^{23} \text{ mol}^{-1}$ and the value of Boltzmann constant is $1.380 \times 10^{-23} \text{ JK}^{-1}$, then the number of significant digits in the calculated value of the universal gas constant is


[JEE(Advanced) 2014]

SOLUTIONS

1. Ans. (0.31 – 0.33)



$$0.02 \qquad \qquad \qquad 0.01$$




$$0.01 \qquad \qquad 0.01$$

$$n_{\text{S}} = 0.01 \text{ mole}$$


$$\text{weight of sulphur} = 32 \times 0.01 = 0.32 \text{ gm}$$

2. Ans. (2.37 - 2.41)

Sol. Mole of $\text{P}_4 = \frac{1.24}{31 \times 4} = 0.01$



$$0.01 \text{ mole} \qquad \qquad \qquad 0.01 \text{ mole}$$



$$0.01 \quad \frac{3}{2} \times 0.01$$

$$= \frac{0.03}{2} \text{ moles}$$

$$W_{\text{CuSO}_4} = \frac{0.03}{2} \times 159 = 2.385 \text{ gm}$$

$$\text{Ans.} = 2.38 \text{ or } 2.39$$

3. Ans. (B, C)

Sol.

Compound	Weight % of P	Weight % of Q
1	50	50
2	44.4	55.6
3	40	60

For option (A)

Let atomic mass of P be M_{P} and atomic mass of Q be M_{Q}

Molar ratio of atoms P : Q in compound 3 is

$$\frac{40}{M_{\text{P}}} : \frac{60}{M_{\text{Q}}} = 3 : 4$$

$$\frac{2M_{\text{Q}}}{3M_{\text{P}}} = \frac{3}{4} \Rightarrow 9M_{\text{P}} = 8M_{\text{Q}}$$

Molar ratio of atoms P : Q in compound 2 is

$$\frac{44.4}{M_{\text{P}}} : \frac{55.6}{M_{\text{Q}}}$$

$$= 44.4 M_{\text{Q}} : 55.6 M_{\text{P}}$$

$$= 44.4 M_{\text{Q}} : 55.6 \times \frac{8M_{\text{Q}}}{9}$$

$$= 44.4 : 55.6 \times \frac{8}{9} = 9 : 10$$

\Rightarrow Empirical formula of compound 2 is therefore P_9Q_{10}

Option (A) is incorrect

So to get 1.29 gm organic salt.

We have to form 0.01 mole salt.

So 0.01 mole nitrobenzene is required.


0.03 mole Sn is required.

So the amount of nitrobenzene = $0.01 \times 123 = 1.23$ gm

the amount of Sn required = $0.01 \times 357 = 3.57$ gm

5. Ans. (1.23)

Sol. The value of y is



So to get 1.29 gm organic salt.

We have to form 0.01 mole salt.

So 0.01 mole nitrobenzene is required.


0.03 mole Sn is required.

So the amount of nitrobenzene = $0.01 \times 123 = 1.23$ gm

the amount of Sn required = $0.01 \times 357 = 3.57$ gm

Ans. 3.57 & 1.23

6. Ans. (6.00 - 6.20)



$$\text{Moles of Al taken} = \frac{5.4}{27} = 0.2$$



$$\text{moles of H}_2\text{SO}_4 \text{ taken} = \frac{50 \times 5.0}{1000} = 0.25$$

$$\text{As } \frac{0.2}{2} > \frac{0.25}{3}, \text{ H}_2\text{SO}_4 \text{ is limiting reagent}$$

$$\text{Now, moles of H}_2 \text{ formed} = \frac{3}{3} \times 0.25 = 0.25$$

$$\therefore \text{Volume of H}_2 \text{ gas formed} = \frac{nRT}{P} = \frac{0.25 \times 0.082 \times 300}{1} = 6.15 \text{ L}$$

7. Ans. (2992)



$$\text{Total mass} = 12 \times 172 + 4 \times 232 = 2992 \text{ g}$$

8. Ans. (4)

Sol. Universal gas constant $R = kN_A$

where k = Boltzman constant and N_A = Avogadro number

$$\begin{aligned} \therefore R &= 1.380 \times 10^{-23} \times 6.023 \times 10^{23} \text{ J/K-mole} \\ &= 8.31174 \\ &\cong 8.312 \end{aligned}$$

So significant figures = 4