

## **DEFINITE INTEGRATION**

1. Let  $f : (0, 1) \rightarrow \mathbb{R}$  be the function defined as  $f(x) = \sqrt{n}$  if  $x \in \left[ \frac{1}{n+1}, \frac{1}{n} \right)$  where  $n \in \mathbb{N}$ . Let  $g : (0, 1) \rightarrow \mathbb{R}$

be a function such that  $\int_{x^2}^x \sqrt{\frac{1-t}{t}} dt < g(x) < 2\sqrt{x}$  for all  $x \in (0, 1)$ . Then  $\lim_{x \rightarrow 0} f(x)g(x)$

[JEE(Advanced) 2023]

2. For  $x \in \mathbb{R}$ , let  $\tan^{-1}(x) \in \left(-\frac{\pi}{2}, \frac{\pi}{2}\right)$ . Then the minimum value of the function  $f : \mathbb{R} \rightarrow \mathbb{R}$  defined by

$$f(x) = \int_0^{x \tan^{-1} x} \frac{e^{(t-\cos t)}}{1+t^{2023}} dt \text{ is}$$

[JEE(Advanced) 2023]

3. Consider the equation

$$\int_1^e \frac{(\log_e x)^{1/2}}{x(a - (\log_e x)^{3/2})^2} dx = 1, \quad a \in (-\infty, 0) \cup (1, \infty).$$

Which of the following statements is/are TRUE ?

[JEE(Advanced) 2022]

- (A) No  $a$  satisfies the above equation
  - (B) An integer  $a$  satisfies the above equation
  - (C) An irrational number  $a$  satisfies the above equation
  - (D) More than one  $a$  satisfy the above equation

4. The greatest integer less than or equal to

$$\int_1^2 \log_2(x^3 + 1) dx + \int_1^{\log_2 9} (2^x - 1)^{\frac{1}{3}} dx$$

is .

[JEE(Advanced) 2022]

5. For positive integer  $n$ , define

$$f(n) = n + \frac{16 + 5n - 3n^2}{4n + 3n^2} + \frac{32 + n - 3n^2}{8n + 3n^2} + \frac{48 - 3n - 3n^2}{12n + 3n^2} + \dots + \frac{25n - 7n^2}{7n^2}.$$

Then, the value of  $\lim_{n \rightarrow \infty} f(n)$  is equal to

[JEE(Advanced) 2022]

- (A)  $3 + \frac{4}{3} \log_e 7$

(B)  $4 - \frac{3}{4} \log_e \left(\frac{7}{3}\right)$

(C)  $4 - \frac{4}{3} \log_e \left(\frac{7}{3}\right)$

(D)  $3 + \frac{3}{4} \log_e 7$

6. Let  $f: \left[-\frac{\pi}{2}, \frac{\pi}{2}\right] \rightarrow \mathbb{R}$  be a continuous function such that

$$f(0) = 1 \text{ and } \int_0^{\frac{\pi}{3}} f(t) dt = 0$$

Then which of the following statements is (are) TRUE?

[JEE(Advanced) 2021]

(A) The equation  $f(x) - 3 \cos 3x = 0$  has at least one solution in  $\left(0, \frac{\pi}{3}\right)$

(B) The equation  $f(x) - 3 \sin 3x = -\frac{6}{\pi}$  has at least one solution in  $\left(0, \frac{\pi}{3}\right)$

(C)  $\lim_{x \rightarrow 0} \frac{x \int_0^x f(t) dt}{1 - e^{x^2}} = -1$

(D)  $\lim_{x \rightarrow 0} \frac{\sin x \int_0^x f(t) dt}{x^2} = -1$

#### Question Stem for Questions Nos. 7 and 8

##### Question Stem

Let  $g_i: \left[\frac{\pi}{8}, \frac{3\pi}{8}\right] \rightarrow \mathbb{R}$ ,  $i = 1, 2$ , and  $f: \left[\frac{\pi}{8}, \frac{3\pi}{8}\right] \rightarrow \mathbb{R}$  be functions such that

$g_1(x) = 1$ ,  $g_2(x) = |4x - \pi|$  and  $f(x) = \sin^2 x$ , for all  $x \in \left[\frac{\pi}{8}, \frac{3\pi}{8}\right]$

Define  $S_i = \int_{\frac{\pi}{8}}^{\frac{3\pi}{8}} f(x) \cdot g_i(x) dx$ ,  $i = 1, 2$

7. The value of  $\frac{16S_1}{\pi}$  is \_\_\_\_.

[JEE(Advanced) 2021]

8. The value of  $\frac{48S_2}{\pi^2}$  is \_\_\_\_.

[JEE(Advanced) 2021]

#### Paragraph for Question No. 9 and 10

Let  $\psi_1: [0, \infty) \rightarrow \mathbb{R}$ ,  $\psi_2: [0, \infty) \rightarrow \mathbb{R}$ ,  $f: [0, \infty) \rightarrow \mathbb{R}$  and  $g: [0, \infty) \rightarrow \mathbb{R}$  be functions such that

$$f(0) = g(0) = 0,$$

$$\psi_1(x) = e^{-x} + x, \quad x \geq 0,$$

$$\psi_2(x) = x^2 - 2x - 2e^{-x} + 2, \quad x \geq 0,$$

$$f(x) = \int_{-x}^x (|t| - t^2) e^{-t^2} dt, \quad x > 0$$

and

$$g(x) = \int_0^{x^2} \sqrt{t} e^{-t} dt, \quad x > 0$$

9. Which of the following statements is **TRUE**? [JEE(Advanced) 2021]

(A)  $f(\sqrt{\ln 3}) + g(\sqrt{\ln 3}) = \frac{1}{3}$

(B) For every  $x > 1$ , there exists an  $\alpha \in (1, x)$  such that  $\psi_1(x) = 1 + \alpha x$

(C) For every  $x > 0$ , there exists a  $\beta \in (0, x)$  such that  $\psi_2(x) = 2x(\psi_1(\beta) - 1)$

(D)  $f$  is an increasing function on the interval  $\left[0, \frac{3}{2}\right]$

10. Which of the following statements is **TRUE**? [JEE(Advanced) 2021]

(A)  $\psi_1(x) \leq 1$ , for all  $x > 0$

(B)  $\psi_2(x) \leq 0$ , for all  $x > 0$

(C)  $f(x) \geq 1 - e^{-x^2} - \frac{2}{3}x^3 + \frac{2}{5}x^5$ , for all  $x \in \left(0, \frac{1}{2}\right)$

(D)  $g(x) \leq \frac{2}{3}x^3 - \frac{2}{5}x^5 + \frac{1}{7}x^7$ , for all  $x \in \left(0, \frac{1}{2}\right)$

11. For any real number  $x$ , let  $[x]$  denote the largest integer less than or equal to  $x$ . If then the value of  $91$  is \_\_\_\_\_. [JEE(Advanced) 2021]

12. Which of the following inequalities is/are **TRUE**? [JEE(Advanced) 2020]

(A)  $\int_0^1 x \cos x dx \geq \frac{3}{8}$

(B)  $\int_0^1 x \sin x dx \geq \frac{3}{10}$

(C)  $\int_0^1 x^2 \cos x dx \geq \frac{1}{2}$

(D)  $\int_0^1 x^2 \sin x dx \geq \frac{2}{9}$

13. Let  $f : \mathbb{R} \rightarrow \mathbb{R}$  be a differentiable function such that its derivative  $f'$  is continuous and  $f(\pi) = -6$ .

If  $F : [0, \pi] \rightarrow \mathbb{R}$  is defined by  $F(x) = \int_0^x f(t) dt$ , and if

$$\int_0^\pi (f'(x) + F(x)) \cos x dx = 2,$$

then the value of  $f(0)$  is \_\_\_\_\_. [JEE(Advanced) 2020]

14. If  $I = \frac{2}{\pi} \int_{-\pi/4}^{\pi/4} \frac{dx}{(1 + e^{\sin x})(2 - \cos 2x)}$  then  $27I^2$  equals \_\_\_\_\_. [JEE(Advanced) 2019]

15. For  $a \in \mathbb{R}$ ,  $|a| > 1$ , let  $\lim_{n \rightarrow \infty} \left( \frac{1 + \sqrt[3]{2} + \dots + \sqrt[3]{n}}{n^{7/3} \left( \frac{1}{(an+1)^2} + \frac{1}{(an+2)^2} + \dots + \frac{1}{(an+n)^2} \right)} \right) = 54$ . Then the possible value(s) of  $a$  is/are :

(A) 8 (B) -9 (C) -6 (D) 7 [JEE(Advanced) 2019]

16. The value of the integral  $\int_0^{\pi/2} \frac{3\sqrt{\cos \theta}}{\left(\sqrt{\cos \theta} + \sqrt{\sin \theta}\right)^5} d\theta$  equals [JEE(Advanced) 2019]

17. For each positive integer  $n$ , let  $y_n = \frac{1}{n}(n+1)(n+2)\dots(n+n)^{1/n}$

For  $x \in \mathbb{R}$ , let  $[x]$  be the greatest integer less than or equal to  $x$ . If  $\lim_{n \rightarrow \infty} y_n = L$ , then the value of  $[L]$  is \_\_\_\_\_.

[JEE(Advanced) 2018]

18. The value of the integral

$$\int_0^{\frac{1}{2}} \frac{1+\sqrt{3}}{((x+1)^2(1-x)^6)^{\frac{1}{4}}} dx$$

is \_\_\_\_\_.

[JEE(Advanced) 2018]

19. Let  $f: \mathbb{R} \rightarrow \mathbb{R}$  be a differentiable function such that  $f(0) = 0$ ,  $f\left(\frac{\pi}{2}\right) = 3$  and  $f'(0) = 1$ . If

$$g(x) = \int_x^{\frac{\pi}{2}} [f'(t) \operatorname{cosec} t - \cot t \operatorname{cosec} t f(t)] dt$$

for  $x \in \left(0, \frac{\pi}{2}\right)$ , then  $\lim_{x \rightarrow 0} g(x) =$

[JEE(Advanced) 2017]

20. If  $I = \sum_{k=1}^{98} \int_k^{k+1} \frac{k+1}{x(x+1)} dx$ , then

[JEE(Advanced)-2017]

(A)  $I < \frac{49}{50}$       (B)  $I < \log_e 99$       (C)  $I > \frac{49}{50}$       (D)  $I > \log_e 99$

21. If  $g(x) = \int_{\sin x}^{\sin(2x)} \sin^{-1}(t) dt$ , then

[JEE(Advanced) 2017]

(A)  $g'\left(\frac{\pi}{2}\right) = -2\pi$       (B)  $g'\left(-\frac{\pi}{2}\right) = 2\pi$   
 (C)  $g'\left(\frac{\pi}{2}\right) = 2\pi$       (D)  $g'\left(-\frac{\pi}{2}\right) = -2\pi$

22. The value of  $\int_{-\frac{\pi}{2}}^{\frac{\pi}{2}} \frac{x^2 \cos x}{1 + e^x} dx$  is equal to

[JEE(Advanced) 2016]

(A)  $\frac{\pi^2}{4} - 2$       (B)  $\frac{\pi^2}{4} + 2$       (C)  $\pi^2 - e^{\frac{\pi}{2}}$       (D)  $\pi^2 + e^{\frac{\pi}{2}}$

23. Let  $f(x) = \lim_{n \rightarrow \infty} \left( \frac{n^n (x+n) \left( x + \frac{n}{2} \right) \dots \left( x + \frac{n}{n} \right)}{n! (x^2 + n^2) \left( x^2 + \frac{n^2}{4} \right) \dots \left( x^2 + \frac{n^2}{n^2} \right)} \right)^{x/n}$ , for all  $x > 0$ . Then

[JEE(Advanced) 2016]

(A)  $f\left(\frac{1}{2}\right) \geq f(1)$       (B)  $f\left(\frac{1}{3}\right) \leq f\left(\frac{2}{3}\right)$       (C)  $f'(2) \leq 0$       (D)  $\frac{f'(3)}{f(3)} \geq \frac{f'(2)}{f(2)}$

24. The total number of distinct  $x \in [0, 1]$  for which  $\int_0^x \frac{t^2}{1+t^4} dt = 2x - 1$  is

[JEE(Advanced) 2016]

25. Let  $f : \mathbb{R} \rightarrow \mathbb{R}$  be a function defined by  $f(x) = \begin{cases} [x] & , \quad x \leq 2 \\ 0 & , \quad x > 2 \end{cases}$ ,

where  $[x]$  is the greatest integer less than or equal to  $x$ . If  $I = \int_{-1}^2 \frac{xf(x^2)}{2+f(x+1)} dx$ , then the value of

$(4I - 1)$  is

[JEE(Advanced) 2015]

26. If  $\alpha = \int_0^1 \left( e^{9x+3\tan^{-1}x} \right) \left( \frac{12+9x^2}{1+x^2} \right) dx$ , where  $\tan^{-1}x$  takes only principal values, then the value of

$\left( \log_e |1+\alpha| - \frac{3\pi}{4} \right)$  is

[JEE(Advanced) 2015]

27. Let  $f : \mathbb{R} \rightarrow \mathbb{R}$  be a continuous odd function, which vanishes exactly at one point and  $f(1) = \frac{1}{2}$ . Suppose

that  $F(x) = \int_{-1}^x f(t) dt$  for all  $x \in [-1, 2]$  and  $G(x) = \int_{-1}^x t |f(f(t))| dt$  for all  $x \in [-1, 2]$ . If  $\lim_{x \rightarrow 1} \frac{F(x)}{G(x)} = \frac{1}{14}$ ,

then the value of  $f\left(\frac{1}{2}\right)$  is

[JEE(Advanced) 2015]

28. The option(s) with the values of  $a$  and  $L$  that satisfy the following equation is(are)

$$\frac{\int_0^{4\pi} e^t (\sin^6 at + \cos^4 at) dt}{\int_0^\pi e^t (\sin^6 at + \cos^4 at) dt} = L ?$$

[JEE(Advanced) 2015]

(A)  $a = 2, L = \frac{e^{4\pi} - 1}{e^\pi - 1}$

(B)  $a = 2, L = \frac{e^{4\pi} + 1}{e^\pi + 1}$

(C)  $a = 4, L = \frac{e^{4\pi} - 1}{e^\pi - 1}$

(D)  $a = 4, L = \frac{e^{4\pi} + 1}{e^\pi + 1}$

29. Let  $f(x) = 7\tan^8 x + 7\tan^6 x - 3\tan^4 x - 3\tan^2 x$  for all  $x \in \left(-\frac{\pi}{2}, \frac{\pi}{2}\right)$ . Then the correct expression(s) is(are)

[JEE(Advanced) 2015]

(A)  $\int_0^{\pi/4} xf(x) dx = \frac{1}{12}$

(B)  $\int_0^{\pi/4} f(x) dx = 0$

(C)  $\int_0^{\pi/4} xf(x) dx = \frac{1}{6}$

(D)  $\int_0^{\pi/4} f(x) dx = 1$

30. Let  $f'(x) = \frac{192x^3}{2 + \sin^4 \pi x}$  for all  $x \in \mathbb{R}$  with  $f\left(\frac{1}{2}\right) = 0$ . If  $m \leq \int_{1/2}^1 f(x)dx \leq M$ , then the possible values of  $m$  and  $M$  are

- (A)  $m = 13, M = 24$       (B)  $m = \frac{1}{4}, M = \frac{1}{2}$   
 (C)  $m = -11, M = 0$       (D)  $m = 1, M = 12$

[JEE(Advanced) 2015]

**Paragraph For Questions 31 and 32**

Let  $F : \mathbb{R} \rightarrow \mathbb{R}$  be a thrice differentiable function. Suppose that  $F(1) = 0, F(3) = -4, F'(x) < 0$  for all  $x \in (1/2, 3)$ . Let  $f(x) = xF(x)$  for all  $x \in \mathbb{R}$ .

31. The correct statement(s) is(are) [JEE(Advanced) 2015]  
 (A)  $f'(1) < 0$       (B)  $f(2) < 0$   
 (C)  $f'(x) \neq 0$  for any  $x \in (1, 3)$       (D)  $f'(x) = 0$  for some  $x \in (1, 3)$

32. If  $\int_1^3 x^2 F'(x)dx = -12$  and  $\int_1^3 x^3 F''(x)dx = 40$ , then the correct expression(s) is(are) [JEE(Advanced) 2015]

- (A)  $9f'(3) + f'(1) - 32 = 0$       (B)  $\int_1^3 f(x)dx = 12$   
 (C)  $9f'(3) - f'(1) + 32 = 0$       (D)  $\int_1^3 f(x)dx = -12$

33. Let  $f : [a, b] \rightarrow [1, \infty)$  be a continuous function and let  $g : \mathbb{R} \rightarrow \mathbb{R}$  be defined as [JEE(Advanced) 2014]

$$g(x) = \begin{cases} 0 & \text{if } x < a \\ \int_a^x f(t)dt & \text{if } a \leq x \leq b \\ \int_a^b f(t)dt & \text{if } x > b \end{cases}$$

Then

- (A)  $g(x)$  is continuous but not differentiable at  $a$   
 (B)  $g(x)$  is differentiable on  $\mathbb{R}$   
 (C)  $g(x)$  is continuous but not differentiable at  $b$   
 (D)  $g(x)$  is continuous and differentiable at either  $a$  or  $b$  but not both.

34. Let  $f : (0, \infty) \rightarrow \mathbb{R}$  be given by  $f(x) = \int_x^1 e^{-\left(\frac{t+1}{t}\right)} \frac{dt}{t}$ . Then [JEE(Advanced) 2014]

- (A)  $f(x)$  is monotonically increasing on  $[1, \infty)$   
 (B)  $f(x)$  is monotonically decreasing on  $[0, 1)$   
 (C)  $f(x) + f\left(\frac{1}{x}\right) = 0$ , for all  $x \in (0, \infty)$   
 (D)  $f(2^x)$  is an odd function of  $x$  on  $\mathbb{R}$

35. The value of  $\int_0^1 4x^3 \left\{ \frac{d^2}{dx^2} (1-x^2)^5 \right\} dx$  is [JEE(Advanced) 2014]

36. The following integral  $\int_{\frac{\pi}{4}}^{\frac{\pi}{2}} (2 \operatorname{cosec} x)^{17} dx$  is equal to - [JEE(Advanced) 2014]

(A)  $\int_0^{\log(1+\sqrt{2})} 2(e^u + e^{-u})^{16} du$

(B)  $\int_0^{\log(1+\sqrt{2})} (e^u + e^{-u})^{17} du$

(C)  $\int_0^{\log(1+\sqrt{2})} (e^u - e^{-u})^{17} du$

(D)  $\int_0^{\log(1+\sqrt{2})} 2(e^u - e^{-u})^{16} du$

37. Let  $f : [0, 2] \rightarrow \mathbb{R}$  be a function which is continuous on  $[0, 2]$  and is differentiable on  $(0, 2)$  with

$f(0) = 1$ . Let  $F(x) = \int_0^{x^2} f(\sqrt{t}) dt$  for  $x \in [0, 2]$ . If  $F'(x) = f'(x)$  for all  $x \in (0, 2)$ , then  $F(2)$  equals -

[JEE(Advanced) 2014]

(A)  $e^2 - 1$

(B)  $e^4 - 1$

(C)  $e - 1$

(D)  $e^4$

### Paragraph For Questions Nos. 38 and 39

Given that for each  $a \in (0, 1)$ ,  $\lim_{h \rightarrow 0^+} \int_h^{1-h} t^{-a} (1-t)^{a-1} dt$  exists. Let this limit be  $g(a)$ . In addition, it is given

that the function  $g(a)$  is differentiable on  $(0, 1)$ .

38. The value of  $g\left(\frac{1}{2}\right)$  is - [JEE(Advanced) 2014]

(A)  $\pi$

(B)  $2\pi$

(C)  $\frac{\pi}{2}$

(D)  $\frac{\pi}{4}$

39. The value of  $g\left(\frac{1}{2}\right)$  is - [JEE(Advanced) 2014]

(A)  $\frac{\pi}{2}$

(B)  $\pi$

(C)  $-\frac{\pi}{2}$

(D) 0

**40.****List-I**

- P.** The number of polynomials  $f(x)$  with non-negative integer coefficients of degree  $\leq 2$ , satisfying

$$f(0) = 0 \text{ and } \int_0^1 f(x)dx = 1, \text{ is}$$

- Q.** The number of points in the interval  $[-\sqrt{13}, \sqrt{13}]$  at which  $f(x) = \sin(x^2) + \cos(x^2)$  attains its maximum value, is

**R.**  $\int_{-2}^2 \frac{3x^2}{(1+e^x)} dx$  equals

**S.**  $\frac{\left( \int_{-\frac{1}{2}}^{\frac{1}{2}} \cos 2x \log\left(\frac{1+x}{1-x}\right) dx \right)}{\left( \int_0^{\frac{1}{2}} \cos 2x \log\left(\frac{1+x}{1-x}\right) dx \right)}$  equals

**List-II**

- 1.** 8

- 2.** 2

- 3.** 4

- 4.** 0

**Codes :****[JEE(Advanced) 2014]**

	P	Q	R	S
(A)	3	2	4	1
(B)	2	3	4	1
(C)	3	2	1	4
(D)	2	3	1	4

## SOLUTIONS

**1. Ans. (C)**

**Sol.**  $\int_{x^2}^x \sqrt{\frac{1-t}{t}} dt \cdot \sqrt{n} \leq f(x)g(x) \leq 2\sqrt{x}\sqrt{n}$

$$\therefore \int_{x^2}^x \sqrt{\frac{1-t}{t}} dt = \sin^{-1} \sqrt{x} + \sqrt{x} \sqrt{1-x} - \sin^{-1} x - x \sqrt{1-x^2}$$

$$\Rightarrow \lim_{x \rightarrow 0} \left( \frac{\sin^{-1} \sqrt{x} + \sqrt{x} \sqrt{1-x} - \sin^{-1} x - x \sqrt{1-x^2}}{\sqrt{x}} \leq f(x)g(x) \leq \frac{2\sqrt{x}}{\sqrt{x}} \right)$$

$$\Rightarrow 2 \leq \lim_{x \rightarrow 0} f(x)g(x) \leq 2$$

$$\Rightarrow \lim_{x \rightarrow 0} f(x)g(x) = 2$$

**2. Ans. (0)**

**Sol.**  $f(x) = \int_0^{x \tan^{-1} x} \frac{e^{t-\cos t}}{1+t^{2023}} dt$


$$f'(x) = \frac{e^{x \tan^{-1} x - \cos(x \tan^{-1} x)}}{1+(x \tan^{-1} x)^{2023}} \cdot \left( \frac{x}{1+x^2} + \tan^{-1} x \right)$$

$$\text{For } x < 0, \tan^{-1} x \in \left(-\frac{\pi}{2}, 0\right)$$

$$\text{For } x \geq 0, \tan^{-1} x \in \left[0, \frac{\pi}{2}\right)$$

$$\Rightarrow x \tan^{-1} x \geq 0 \quad \forall x \in \mathbb{R}$$

$$\text{And } \frac{x}{1+x^2} + \tan^{-1} x = \begin{cases} > 0 & \text{For } x > 0 \\ < 0 & \text{For } x < 0 \\ 0 & \text{For } x = 0 \end{cases}$$



$$\text{Hence minimum value is } f(0) = \int_0^0 = 0$$

**3. Ans. (C, D)**

**Sol.**  $\int_1^e \frac{(\log_e x)^{1/2}}{x \left( a - (\log_e x)^{3/2} \right)^2} dx = 1$

$$\text{Let } a - (\log_e x)^{3/2} = t$$

$$\frac{(\log_e x)^{1/2}}{x} dx = -\frac{2}{3} dt$$

$$= \frac{2}{3} \int_a^{a-1} \frac{-dt}{t^2} = \frac{2}{3} \left( \frac{1}{t} \right)_a^{a-1} = 1$$

$$\frac{2}{3a(a-1)} = 1$$

$$3a^2 - 3a - 2 = 0$$

$$a = \frac{3 \pm \sqrt{33}}{6}$$

**4. Ans. (5)**

**Sol.**  $f(x) = \log_2(x^3 + 1) = y$

$$x^3 + 1 = 2^y \Rightarrow x = (2^y - 1)^{1/3} = f^{-1}(y)$$

$$f^{-1}(x) = (2^x - 1)^{1/3}$$

$$= \int_1^2 \log_2(x^3 + 1) dx + \int_1^{\log_2 9} (2^x - 1)^{1/3} dx$$

$$= \int_1^2 f(x) dx + \int_1^{\log_2 9} f^{-1}(x) dx = 2 \log_2 9 - 1$$

$$= 8 < 9 < 2^{7/2} \Rightarrow 3 < \log_2 9 < \frac{7}{2}$$

$$= 5 < 2 \log_2 9 - 1 < 6$$

$$[2 \log_2 9 - 1] = 5$$

**5. Ans. (B)**

**Sol.**  $f(n) = n + \sum_{r=1}^n \frac{16r + (9-4r)n - 3n^2}{4rn + 3n^2}$

$$f(n) = n + \sum_{r=1}^n \frac{(16r + 9n) - (4rn + 3n^2)}{4rn + 3n^2}$$

$$f(n) = n + \left( \sum_{r=1}^n \frac{16r + 9n}{4rn + 3n^2} \right) - n$$

$$\lim_{n \rightarrow \infty} f(n) = \lim_{n \rightarrow \infty} \sum \frac{16r + 9n}{4rn + 3n^2}$$

$$= \lim_{n \rightarrow \infty} \sum_{r=1}^n \frac{\left( 16 \left( \frac{r}{n} \right) + 9 \right) \frac{1}{n}}{4 \left( \frac{r}{n} \right) + 3}$$

$$= \int_0^1 \frac{16x + 9}{4x + 3} dx = \int_0^1 4 dx - \int_0^1 \frac{3}{4x + 3} dx$$

$$= 4 - \frac{3}{4} (\ell n |4x + 3|)_0^1$$

$$= 4 - \frac{3}{4} \ell n \frac{7}{3}$$

**6. Ans. (A, B, C)**

**Sol.** (A) Let  $g(x) = f(x) - 3 \cos 3x$

Now,

$$\int_0^{\pi/3} g(x)dx = \int_0^{\pi/3} f(x)dx - 3 \int_0^{\pi/3} \cos 3x dx = 0$$

Hence  $g(x) = 0$  has a root in  $\left(0, \frac{\pi}{3}\right)$

$$(B) \text{ Let } h(x) = f(x) - 3 \sin 3x + \frac{6}{\pi}$$

Now,

$$\begin{aligned} \int_0^{\pi/3} h(x)dx &= \int_0^{\pi/3} f(x)dx - 3 \int_0^{\pi/3} \sin 3x dx + \int_0^{\pi/3} \frac{6}{\pi} dx \\ &= 0 - 2 + 2 = 0 \end{aligned}$$

Hence  $h(x) = 0$  has a root in  $\left(0, \frac{\pi}{3}\right)$

$$(C) \lim_{x \rightarrow 0} \frac{x \int_0^x f(t) dt}{1 - e^{x^2}} = \lim_{x \rightarrow 0} \underbrace{\left( \frac{x^2}{1 - e^{x^2}} \right)}_{\substack{\text{Apply L'Hopital's Rule}}} \underbrace{\frac{\int_0^x f(t) dt}{x}}$$

$$= -1 \lim_{x \rightarrow 0} \frac{f(x)}{1} = -1$$

$$(D) \lim_{x \rightarrow 0} \frac{(\sin x) \int_0^x f(t) dt}{x^2}$$

$$= \lim_{x \rightarrow 0} \underbrace{\left( \frac{\sin x}{x} \right)}_1 \underbrace{\frac{\int_0^x f(t) dt}{x}}_{\substack{\text{Apply L'Hopital's Rule}}}$$

$$= 1 \lim_{x \rightarrow 0} \frac{f(x)}{1} = 1$$

**7. Ans. (2.00)**

**Sol.**

$$S_1 = \int_{\pi/8}^{3\pi/8} f(x)dx = \int_{\pi/8}^{3\pi/8} \sin^2 x dx = \int_{\pi/8}^{3\pi/8} \sin^2 \left( \frac{\pi}{8} + \frac{3\pi}{8} - x \right) dx$$


$$= \int_{\pi/8}^{3\pi/8} \cos^2 x dx$$

$$2S_1 = \int_{\pi/8}^{3\pi/8} (\sin^2 x + \cos^2 x) dx = \frac{3\pi}{8} - \frac{\pi}{8} = \frac{\pi}{4}$$

$$\Rightarrow \frac{16S_1}{\pi} = 2$$

**8. Ans. (1.50)**

**Sol.**



$$S_2 = \int_{\pi/8}^{3\pi/8} f(x)g_2(x)dx = \int_{\pi/8}^{3\pi/8} \sin^2 x |4x - \pi| dx$$

$$= \int_{\pi/8}^{3\pi/8} \sin^2 \left( \frac{\pi}{2} - x \right) \left| 4 \left( \frac{\pi}{2} - x \right) - \pi \right| dx$$

$$= \int_{\pi/8}^{3\pi/8} (\cos^2 x) |4x - \pi| dx$$

$$\Rightarrow 2S_2 = \int_{\pi/8}^{3\pi/8} |4x - \pi| (\sin^2 x + \cos^2 x) dx$$

$$= \int_{\pi/8}^{3\pi/8} |4x - \pi| dx$$

$$= 2 \times \frac{1}{2} \times \frac{\pi}{8} \times \frac{\pi}{2} = \frac{\pi^2}{16}$$

$$\Rightarrow \frac{48S_2}{\pi^2} = \frac{3}{2} = 1.5$$

**9. Ans. (C)**

$$\text{Sol. } f'(x) = (|x| - x^2)e^{-x^2} + (|x| - x^2)e^{-x^2}, \quad x \geq 0$$

$$f = 2(x - x^2) e^{-x^2}$$

$$\begin{array}{c} + + + \\ \hline 0 \quad 1 \quad - - - \end{array}$$

hence option (D) is wrong

$$g'(x) = xe^{-x^2} 2x$$

$$f'(x) + g'(x) = 2 \times e^{-x^2}$$

$$f(x) + g(x) = -e^{-x^2} + c$$

$$f(x) + g(x) = -e^{-x^2} + 1$$

$$f(\ln 3) + g(\sqrt{\ln 3}) = 1 - \frac{1}{3} = \frac{2}{3}$$

(option (A) is wrong)

$$H(x) = \psi_1(x) - 1 - \alpha x = e^{-x} + x - 1 - \alpha x,$$

$x \geq 1$  &  $\alpha \in (1, x)$

$$H(1) = e^{-1} + 1 - 1 - \alpha < 0$$

$$H'(x) = -e^{-x} + 1 - \alpha > 0 \Rightarrow H(x) \text{ is } \downarrow$$

$\Rightarrow$  option (B) is wrong

$$(C) \psi_2(x) = 2(\psi_1(\beta) - 1)$$

Applying L.M.V.T to  $\psi_2(x)$  in  $[0, x]$

$$\psi'_2(\beta) = \frac{\psi_2(x) - \psi_2(0)}{x}$$

$$2\beta - 2 + 2e^{-\beta} = \frac{\psi_2(x) - 0}{x}$$

$\Rightarrow \psi_2(x) = 2x(\psi_1(\beta) - 1)$  has one solution

option (C) is correct.

### 10. Ans. (D)

$$\text{Sol. (A)} \quad \psi_1(x) = e^{-x} + x, \quad x \geq 0$$

$$\psi'_1(x) = 1 - e^{-x} > 0 \Rightarrow \psi_1(x) \text{ is } \uparrow$$

$$\psi_1(x) \geq \psi_1(0) \quad \forall x \geq 0 \Rightarrow \psi_1(x) \geq 1$$

$$(B) \quad \psi_2(x) = x^2 - 2x + 2 - 2e^{-x} \quad x \geq 0$$

$$\psi'_2(x) = 2x - 2 + 2e^{-x} = 2\psi_1(x) - 2 \geq 0 \quad \forall x \geq 0$$

$\Rightarrow \psi_2(x) \text{ is } \uparrow$

$$\Rightarrow \psi_2(x) \geq \psi_2(0) \Rightarrow \psi_2(x) \geq 0$$

$$(C) \quad f(x) = 2 \int_0^x (t - t^2) e^{-t^2} dt \quad \& \quad x \in \left(0, \frac{1}{2}\right)$$

$$\text{Let, } H(x) = f(x) - 1 + e^{-x^2} + \frac{2}{3}x^3 - \frac{2}{5}x^5, \quad x \in \left(0, \frac{1}{2}\right)$$

$$H(0) = 0$$

$$H'(x) = 2(x - x^2)e^{-x^2} - 2xe^{-x^2} + 2x^2 - 2x^4$$

$$= -2x^2 e^{-x^2} + 2x^2 - 2x^4$$

$$= 2x^2(1 - x^2 - e^{-x^2})$$

$$\because e^{-x} \geq 1 - x \quad \forall x \geq 0$$

$$\Rightarrow H'(x) \leq 0$$

$$\Rightarrow H(x) \text{ is } \downarrow \Rightarrow H(x) \leq 0 \quad \forall x \in \left(0, \frac{1}{2}\right)$$

$$f(x) \leq 1 - e^{-x^2} - \frac{2}{3}x^3 + \frac{2}{5}x^5; \quad \forall x \in \left(0, \frac{1}{2}\right)$$

$$\text{Let } P(x) = g(x) - \frac{2}{3}x^3 + \frac{2}{5}x^5 - \frac{1}{7}x^7 \quad x \in \left(0, \frac{1}{2}\right)$$

$$P'(x) = 2x^2 e^{-x^2} - 2x^2 + 2x^4 - x^6$$

$$= 2x^2 \left(1 - \frac{x^2}{1} + \frac{x^4}{2} - \frac{x^6}{3} + \dots\right) - 2x^2 + 2x^4 - x^6$$

$$= -\frac{x^8}{3} + \frac{x^{10}}{12} \dots$$

$$\Rightarrow P'(x) \leq 0$$

$$\Rightarrow P(x) \text{ is } \downarrow$$

$$\Rightarrow P(x) \leq 0$$

option (D) is correct

### 11. Ans. (182)

$$\text{Sol. Let } f(x) = \left(\frac{10x}{x+1}\right)$$

$$\text{So, } f'(x) = 10 \left(\frac{(x+1)-x}{(x+1)^2}\right) = \frac{10}{(x+1)^2} > 0$$

$$\forall x \in [0, 10],$$

So,  $f(x)$  is an increasing function

$$\text{So, range of } f(x) \text{ is } \left[0, \sqrt{\frac{100}{11}}\right]$$

$$I = \int_0^{1/9} \left[ \sqrt{\frac{10x}{x+1}} \right] dx + \int_{2/3}^9 \left[ \sqrt{\frac{10x}{x+1}} \right] dx +$$

$$\int_{1/9}^{2/3} \left[ \sqrt{\frac{10x}{x+1}} \right] dx + \int_{2/3}^9 \left[ \sqrt{\frac{10x}{x+1}} \right] dx + \int_9^{10} \left[ \sqrt{\frac{10x}{x+1}} \right] dx$$

$$= 0 + \int_{1/9}^{2/3} dx + 2 \int_{2/3}^9 dx + 3 \int_9^{10} dx$$

$$= \frac{2}{3} - \frac{1}{9} + 2 \left(9 - \frac{2}{3}\right) + 3(10 - 9)$$

$$= \frac{6-1}{9} + 2 \times \frac{25}{3} + 3 = \frac{5}{9} + \frac{50}{3} + 3$$

$$= \frac{5+150+27}{9} = \frac{182}{9} \Rightarrow 9I = 182$$

### 12. Ans. (A, B, D)

$$\text{Sol. (A)} \quad \cos x = 1 - \frac{x^2}{2!} + \frac{x^4}{4!} - \dots$$

$$\sin x = x - \frac{x^3}{3!} + \frac{x^5}{5!} - \dots$$

$$\cos x \geq 1 - \frac{x^2}{2}$$

$$\int_0^1 x \cos x \geq \int_0^1 x \left(1 - \frac{x^2}{2}\right) = \frac{1}{2} - \frac{1}{8}$$

$$\int_0^1 x \cos x \geq \frac{3}{8} \quad (\text{True})$$

(B)  $\sin x \geq x - \frac{x^3}{6}$

$$\int_0^1 x \sin x \geq \int_0^1 x \left(x - \frac{x^3}{6}\right) dx$$

$$\int_0^1 x \sin x \geq \frac{1}{3} - \frac{1}{30} \Rightarrow \int_0^1 x \sin x dx \geq \frac{3}{10}$$

(True)

(D)  $\int_0^1 x^2 \sin x dx \geq \int_0^1 x^2 \left(x - \frac{x^3}{6}\right) dx$

$$\int_0^1 x^2 \sin x dx \geq \frac{1}{4} - \frac{1}{36}$$

$$\int_0^1 x^2 \sin x dx \geq \frac{2}{9} \quad (\text{True})$$

(C)  $\cos x < 1$

$$x^2 \cos x < x^2$$

$$\int_0^1 x^2 \cos x dx < \int_0^1 x^2 dx$$

$$\int_0^1 x^2 \cos x dx < \frac{1}{3}$$

So, option (C) is incorrect.

**13. Ans. (1080.00)**

**Sol.**  $F(x) = \int_0^x f(t) dt$

$$\Rightarrow F'(x) = f(x)$$

$$I = \int_0^\pi f'(x) \cos x dx + \int_0^\pi F(x) \cos(x) dx = 2 \dots (1)$$

$$I_1 = \int_0^\pi f'(x) \cos x dx \quad (\text{Let})$$

Using by parts

$$I_1 = (\cos x \cdot f(x))_0^\pi + \int_0^\pi \sin x \cdot f(x) dx$$

$$I_1 = 6 - f(0) + \int_0^\pi \sin x \cdot F'(x) dx$$

$$I_1 = 6 - f(0) + I_2 \dots (2)$$

$$I_2 = \int_0^\pi \sin x \cdot F'(x) dx$$

Using by part we get

$$I_2 = (\sin x \cdot F(x))_0^\pi - \int_0^\pi \cos x \cdot F(x) dx$$

$$I_2 = - \int_0^\pi \cos x \cdot F(x) dx$$

$$(2) \Rightarrow I_1 = 6 - f(0) - \int_0^\pi \cos x \cdot F(x) dx$$

$$(1) \Rightarrow I = 6 - f(0) = 2 \Rightarrow f(0) = 4$$

**14. Ans. (4.00)**

**Sol.**

$$2I = \frac{2}{\pi} \int_{-\pi/4}^{\pi/4} \left[ \frac{1}{(1+e^{\sin x})(2-\cos 2x)} + \frac{1}{(1+e^{-\sin x})(2-\cos 2x)} \right] dx$$

(using King's Rule)

$$\Rightarrow I = \frac{1}{\pi} \int_{-\pi/4}^{\pi/4} \frac{dx}{2-\cos 2x}$$

$$\Rightarrow I = \frac{2}{\pi} \int_0^{\pi/4} \frac{dx}{2-\cos 2x} = \frac{2}{\pi} \int_0^{\pi/4} \frac{\sec^2 x dx}{1+3\tan^2 x}$$

$$= \frac{2}{\sqrt{3}\pi} \left[ \tan^{-1}(\sqrt{3} \tan x) \right]_0^{\pi/4} = \frac{2}{3\sqrt{3}}$$

$$\Rightarrow 27I^2 = 27 \times \frac{4}{27} = 4$$

**15. Ans. (A, B)**

**Sol.**  $\lim_{n \rightarrow \infty} \frac{n^{1/3} \left( \sum_{r=1}^n \left( \frac{r}{n} \right)^{1/3} \right)}{n^{7/3} \left( \sum_{r=1}^n \frac{1}{(an+r)^2} \right)} = 54$

$$\lim_{n \rightarrow \infty} \left( \frac{\frac{1}{n} \sum_{r=1}^n \left( \frac{r}{n} \right)^{1/3}}{\frac{1}{n} \sum_{r=1}^n \frac{1}{(a+r/n)^2}} \right) = 54$$

$$\begin{aligned} & \int_0^1 x^{1/3} dx \\ \Rightarrow & \frac{\int_0^1 \frac{1}{(a+x)^2} dx}{\int_0^1 \frac{1}{(a+x)^2} dx} = 54 \\ & \Rightarrow \frac{\frac{3}{4}}{\frac{1}{a(a+1)}} = 54 \\ & \Rightarrow a(a+1) = 72 \\ & \Rightarrow a^2 + a - 72 = 0 \Rightarrow a = -9, 8 \end{aligned}$$

16. Ans. (0.50)

$$\begin{aligned} \text{Sol. } I &= \int_0^{\pi/2} \frac{3\sqrt{\cos \theta}}{\left(\sqrt{\cos \theta} + \sqrt{\sin \theta}\right)^5} d\theta \\ &= \int_0^{\pi/2} \frac{3\sqrt{\sin \theta}}{\left(\sqrt{\cos \theta} + \sqrt{\sin \theta}\right)^5} d\theta \\ 2I &= \int_0^{\pi/2} \frac{3d\theta}{\left(\sqrt{\cos \theta} + \sqrt{\sin \theta}\right)^4} \\ &= 3 \int_0^{\pi/2} \frac{\sec^2 \theta d\theta}{\left(1 + \sqrt{\tan \theta}\right)^4} \end{aligned}$$

$$\text{Let } 1 + \sqrt{\tan \theta} = t$$

$$\begin{aligned} \frac{\sec^2 \theta}{2\sqrt{\tan \theta}} d\theta &= dt \\ \sec^2 \theta d\theta &= 2(t-1)dt \end{aligned}$$

$$= 3 \int_1^\infty \frac{2(t-1)dt}{t^4}$$

$$= 6 \int_1^\infty (t^{-3} - t^{-4}) dt$$

$$2I = 6 \left( \frac{t^{-2}}{-2} - \frac{t^{-3}}{-3} \right)_1^\infty = 6 \left[ 0 - 0 - \left\{ -\frac{1}{2} + \frac{1}{3} \right\} \right]$$

$$I = 0.50$$

17. Ans. (1)

$$\begin{aligned} \text{Sol. } y_n &= \left\{ \left( 1 + \frac{1}{n} \right) \left( 1 + \frac{2}{n} \right) \dots \left( 1 + \frac{n}{n} \right) \right\}^{\frac{1}{n}} \\ y_n &= \prod_{r=1}^n \left( 1 + \frac{r}{n} \right)^{1/n} \end{aligned}$$

$$\begin{aligned} \log y_n &= \frac{1}{n} \sum_{r=1}^n \ell n \left( 1 + \frac{r}{n} \right) \\ \Rightarrow \lim_{n \rightarrow \infty} \log y_n &= \lim_{x \rightarrow \infty} \sum_{r=1}^n \frac{1}{n} \ell n \left( 1 + \frac{r}{n} \right) \\ \Rightarrow \log L &= \int_0^1 \ell n(1+x) dx \\ \Rightarrow \log L &= \log \frac{4}{e} \\ \Rightarrow L &= \frac{4}{e} \\ \Rightarrow [L] &= 1 \end{aligned}$$

18. Ans. (2)

$$\begin{aligned} \text{Sol. } & \int_0^{\frac{1}{2}} \frac{(1+\sqrt{3})dx}{\left[(1+x)^2(1-x)^6\right]^{1/4}} \\ & \int_0^{\frac{1}{2}} \frac{(1+\sqrt{3})dx}{(1+x)^2 \left[ \frac{(1-x)^6}{(1+x)^6} \right]^{1/4}} \\ \text{Put } \frac{1-x}{1+x} &= t \Rightarrow = \frac{-2dx}{(1+x)^2} dt \\ I &= \int_1^{\frac{1}{2}} \frac{(1+\sqrt{3})dt}{-2t^{6/4}} = \frac{-(1+\sqrt{3})}{2} \times \left| \frac{-2}{\sqrt{t}} \right|_1^{\frac{1}{2}} \\ &= (1+\sqrt{3})(\sqrt{3}-1) = 2 \end{aligned}$$

19. Ans. (2)

$$\begin{aligned} \text{Sol. } g(x) &= \int_x^{\pi/2} (f'(t) \operatorname{cosec} t - f(t) \operatorname{cosec} t \cot t) dt \\ &= \int_x^{\pi/2} (f(t) \operatorname{cosec} t)' dt \\ &= f\left(\frac{\pi}{2}\right) \operatorname{cosec}\left(\frac{\pi}{2}\right) - \frac{f(x)}{\sin x} = 3 - \frac{f(x)}{\sin x} \\ \therefore \lim_{x \rightarrow 0} g(x) &= 3 - \lim_{x \rightarrow 0} \frac{f(x)}{\sin x}; \text{ as } f'(0) = 1 \\ \Rightarrow \lim_{x \rightarrow 0} g(x) &= 3 - 1 = 2 \end{aligned}$$

**20. Ans. (B,C)**

$$\text{Sol. } S_k = \sum_{k=1}^{98} \int_k^{k+1} \frac{k+1}{x(x+1)} dx = \sum_{k=1}^{98} I_k$$

$$1 \leq k \leq x \leq k+1 \leq 99$$

$$2 \leq k+1 \leq x+1 \leq k+2 \leq 100$$

$$\frac{k+1}{(k+1)(100)} \leq \frac{k+1}{x(x+1)} \leq \frac{k+1}{x(x+1)}$$

$$\int_k^{k+1} \frac{1}{100} dx \leq \int_k^{k+1} \frac{k+1}{x(x+1)} dx \leq \int_k^{k+1} \frac{1}{x} dx$$

$$\frac{1}{100} \leq I_k \leq \ln\left(\frac{k+1}{k}\right)$$

$$\frac{98}{100} \leq \sum_{k=1}^{98} I_k \leq \ln 99$$

$$\frac{49}{50} \leq S_k \leq \ln 99$$

**Aliter**

$$\begin{aligned} I &= \sum_{k=1}^{98} \left( \int_k^{k+1} \frac{(k+1)}{x(x+1)} dx \right) \\ &= \sum_{k=1}^{98} (k+1) \left( \int_k^{k+1} \left( \frac{1}{x} - \frac{1}{x+1} \right) dx \right) \\ &= \sum_{k=1}^{98} (k+1) \left( (\ln x - \ln(x+1))_k^{k+1} \right) \\ &= \sum_{k=1}^{98} (k+1) (\ln(k+1) - \ln(k+2) - \ln k + \ln(k+1)) \\ &= \sum_{k=1}^{98} ((k+1) \ln(k+1) - k \ln k) - \\ &\quad \sum_{k=1}^{98} ((k+1) \ln(k+2) - k \ln(k+1)) + \sum_{k=1}^{98} (\ln(k+1) - \ln k) \\ &\quad \text{(Difference series)} \\ \therefore I &= (99 \ln 99) + (-99 \ln 100 + \ln 2) + (\ln 99) \\ &= \ln \left( \frac{2 \times (99)^{100}}{(100)^{99}} \right) \quad \dots\dots(1) \end{aligned}$$

For option (B) :

$$\begin{aligned} \text{Now, consider } (100)^{99} &= (1+99)^{99} \\ &= {}^{99}C_0 + {}^{99}C_1(99) + {}^{99}C_2(99)^2 + \dots + \\ &\quad {}^{99}C_{97}(99)^{97} + \underbrace{{}^{99}C_{98}(99)^{98}}_{\text{(value}=(99)^{99})} + \underbrace{{}^{99}C_{99}(99)^{99}}_{\text{(value}=(99)^{99})} \end{aligned}$$

$$\Rightarrow (100)^{99} > 2.(99)^{99} \Rightarrow \frac{2 \times (99)^{99}}{(100)^{99}} < 1$$

$$\therefore \frac{2 \times (99)^{100}}{(100)^{99}} < 99 \text{ (on multiplying by 99)}$$

$$\Rightarrow I < \ln 99$$

For option (C) :

$$\text{Since, } \sum_{k=1}^{98} \int_k^{k+1} \frac{k+1}{(x+1)^2} dx < \sum_{k=1}^{98} \int_k^{k+1} \frac{(k+1)dx}{x(x+1)}$$

$$\Rightarrow \sum_{k=1}^{98} \left( \frac{1}{k+2} \right) < I$$

(on integration)

$$\Rightarrow \underbrace{\left( \frac{1}{3} + \frac{1}{4} + \frac{1}{5} + \dots + \frac{1}{100} \right)}_{98 \text{ terms}} < I$$

$$\Rightarrow \frac{98}{100} < \frac{1}{3} + \frac{1}{4} + \frac{1}{5} + \dots + \frac{1}{100} < I$$

$$\therefore I > \frac{49}{50}$$

Hence option (C) is correct.

**21. Ans. (BONUS)**

$$\text{Sol. } g(x) = \int_{\sin x}^{\sin 2x} \sin^{-1} t dt$$

$$\Rightarrow g'(x) = 2\sin^{-1}(\sin 2x) \times \cos 2x - \sin^{-1}(\sin x) \cos x$$

$$\Rightarrow g'\left(\frac{\pi}{2}\right) = 0 \text{ & } g'\left(-\frac{\pi}{2}\right) = 0$$

No option matches the result

**22. Ans. (A)**

$$\text{Sol. Let } I = \int_{-\pi/2}^{\pi/2} \frac{x^2 \cos x}{1+e^x} dx$$

$$= \int_0^{\pi/2} \left( \frac{1}{1+e^x} + \frac{1}{1+e^{-x}} \right) x^2 \cos x dx$$

$$= \int_0^{\pi/2} x^2 \cos x dx = \left( x^2 \sin x \right)_0^{\pi/2} - 2 \int_0^{\pi/2} x \cdot \sin x dx \quad \text{(I) (II)}$$

$$= \frac{\pi^2}{4} - 2 \left[ - \left( x \cdot \cos x \right)_0^{\pi/2} + \int_0^{\pi/2} 1 \cdot \cos x dx \right]$$

$$= \frac{\pi^2}{4} - 2[0+1] = \left( \frac{\pi^2}{4} - 2 \right)$$

**23. Ans. (B, C)**

**Sol.**

$$\ln f(x) = \lim_{n \rightarrow \infty} \frac{x}{n} \ln \left[ \frac{\prod_{r=1}^n \left( x + \frac{1}{r/n} \right)}{\prod_{r=1}^n \left( x^2 + \frac{1}{(r/n)^2} \right)} \frac{1}{\prod_{r=1}^n (r/n)} \right]$$

$$= x \lim_{n \rightarrow \infty} \frac{1}{n} \sum_{r=1}^n \ln \left( \frac{x \frac{r}{n} + 1}{\left( x \frac{r}{n} \right)^2 + 1} \right)$$

$$= x \int_0^1 \ln \left( \frac{1+tx}{1+t^2x^2} \right) dt \quad (\text{put } tx = z)$$

$$\ln f(x) = \int_0^x \ln \left( \frac{1+z}{1+z^2} \right) dz$$

$$\Rightarrow \frac{f'(x)}{f(x)} = \ln \left( \frac{1+x}{1+x^2} \right)$$

sign scheme of  $f'(x)$  

also  $f'(1) = 0$

$$\Rightarrow f\left(\frac{1}{2}\right) < f(1), f\left(\frac{1}{3}\right) < f\left(\frac{2}{3}\right), f'(2) < 0$$

$$\text{Also } \frac{f'(3)}{f(3)} - \frac{f'(2)}{f(2)} = \ln \left( \frac{4}{10} \right) - \ln \left( \frac{3}{5} \right)$$

$$= \ln \left( \frac{4}{6} \right) < 0 \Rightarrow \frac{f'(3)}{f(3)} < \frac{f'(2)}{f(2)}$$

**24. Ans. (1)**

$$\text{Sol. Let } f(x) = \int_0^x \frac{t^2}{1+t^4} dt - 2x + 1$$

$$\Rightarrow f'(x) = \frac{x^2}{1+x^4} - 2$$

$$\text{as } \frac{1+x^4}{x^2} \geq 2 \Rightarrow \frac{x^2}{1+x^4} \leq \frac{1}{2}$$

$$\Rightarrow f'(x) \leq -\frac{3}{2} \Rightarrow f(x) \text{ is continuous and decreasing}$$

$$f(0) = 1 \text{ and } f(1) = \int_0^1 \frac{t^2}{1+t^4} dt - 2 \leq -\frac{3}{2}$$

by IVT  $f(x) = 0$  possesses exactly one solution in  $[0, 1]$

**25. Ans. (0)**

$$\text{Sol. Given } f(x) = \begin{cases} [x] & x \leq 2 \\ 0 & x > 2 \end{cases}$$

where  $[x]$  denotes greatest integer function.

$$\text{Now } I = \int_{-1}^2 \frac{xf(x^2)}{2+f(x+1)} dx$$

$$I = \int_{-1}^0 \frac{xf(x^2)}{2+f(x+1)} dx + \int_0^1 \frac{xf(x^2)}{2+f(x+1)} dx + \int_1^{\sqrt{2}} \frac{xf(x^2)}{2+f(x+1)} dx$$

$$+ \int_{\sqrt{2}}^{\sqrt{3}} \frac{xf(x^2)}{2+f(x+1)} dx + \int_{\sqrt{3}}^2 \frac{xf(x^2)}{2+f(x+1)} dx$$

$$\therefore I = I_1 + I_2 + I_3 + I_4 + I_5$$

Clearly  $I_1, I_2, I_4$  &  $I_5$  are zero using definition of  $f(x)$

$$\therefore I = I_3 = \int_1^{\sqrt{2}} \frac{xf(x^2)}{2+f(x+1)} dx$$

$$= \int_1^{\sqrt{2}} \frac{x \cdot 1}{2+0} dx = \frac{x^2}{4} \Big|_1^{\sqrt{2}} = \frac{2}{4} - \frac{1}{4} = \frac{1}{4}$$

$$\therefore 4I - 1 = 0$$

**26. Ans. (9)**

$$\text{Sol. } \alpha = \int_0^1 e^{(9x+3\tan^{-1}x)} \left( 9 + \frac{3}{1+x^2} \right) dx$$

$$\alpha = \left( e^{9x+3\tan^{-1}x} \right)_0^1$$

$$= e^{9+\frac{3\pi}{4}} - 1$$

$$\log |\alpha + 1| = 9 + \frac{3\pi}{4} \Rightarrow \text{Ans.} = 9$$

**27. Ans. (7)**

$$\text{Sol. } \lim_{x \rightarrow 1} \frac{F(x)}{G(x)} = \lim_{x \rightarrow 1} \frac{F'(x)}{G'(x)}$$

$$= \lim_{x \rightarrow 1} \frac{f(x)}{x |f(f(x))|} = \frac{1}{14}$$

$$\Rightarrow \frac{\frac{1}{2}}{|f\left(\frac{1}{2}\right)|} = \frac{1}{14} \Rightarrow \left| f\left(\frac{1}{2}\right) \right| = 7$$

$$f\left(\frac{1}{2}\right) = 7$$

$$f\left(\frac{1}{2}\right) \neq -7 \text{ as } f(x) \text{ vanishes exactly at one point.}$$

**28. Ans. (A, C)**

**Sol.** Let  $I_1 = \int_0^{4\pi} e^t (\sin^6 at + \cos^4 at) dt$

$$= \int_0^\pi e^t (\sin^6 at + \cos^4 at) dt + \int_\pi^{2\pi} e^t (\sin^6 at + \cos^4 at) dt$$

$$+ \int_{2\pi}^{3\pi} e^t (\sin^6 at + \cos^6 at) dt + \int_{3\pi}^{4\pi} e^t (\sin^6 at + \cos^6 at) dt$$

$$\therefore I_1 = \int_0^\pi e^t (\sin^6 at + \cos^4 at) dt +$$

$$\int_0^{\pi+t} e^{\pi+t} (\sin^6 at + \cos^4 at) dt +$$

$$+ \int_0^{\pi+3\pi} e^{1+3\pi} (\sin^6 at + \cos^4 at) dt$$

$$= (1 + e^\pi + e^{2\pi} + e^{3\pi}) \int_0^\pi e^t (\sin^6 at + \cos^4 at) dt$$

$$\frac{\int_0^\pi e^t (\sin^6 at + \cos^4 at) dt}{\int_0^\pi e^t (\sin^6 at + \cos^4 at) dt} = 1 + e^\pi + e^{2\pi} + e^{3\pi} = \frac{e^{4\pi} - 1}{e^\pi - 1}$$

**29. Ans. (A, B)**

**Sol.** Given  $f(x) = (7 \tan^6 x - 3 \tan^2 x) \sec^2 x$

$$\therefore \int_0^{\pi/4} \underbrace{x \left( 7 \tan^6 x - 3 \tan^2 x \right) \sec^2 x dx}_{\text{II}}$$

Using I.B.P.

$$= x \left( \tan^7 x - \tan^3 x \right)_0^{\pi/4} - \int_0^{\pi/4} (\tan^7 x - \tan^3 x) dx$$

$$= - \int_0^{\pi/4} \tan^3 x (\tan^2 x - 1) \sec^2 x dx$$

Put  $\tan x = t$ 

$$= \int_0^1 (t^3 - t^5) dt = \frac{1}{4} - \frac{1}{6} = \frac{3-2}{12} = \frac{1}{12}$$

Also,

$$\int_0^{\pi/4} (7 \tan^6 x - 3 \tan^2 x) \sec^2 x dx$$

$$= \tan^7 x - \tan^3 x \Big|_0^{\pi/4} = 0$$

**30. Ans. (D)**

**Sol.**  $f(x) = \frac{192x^3}{2 + \sin^4 \pi x}$

$$\frac{192x^3}{3} \leq f'(x) \leq \frac{192x^3}{2}$$

$$\frac{192}{12} \left( x^4 - \frac{1}{16} \right) \leq \int_{1/2}^x f'(x) dx \leq \frac{192}{8} \left( x^4 - \frac{1}{16} \right)$$

$$16x^4 - 1 \leq f(x) \leq 24x^4 - \frac{3}{2}$$

$$2.6 \leq \int_{1/2}^1 f(x) dx \leq 3.9$$

out of given options only option (D) is correct.

**31. Ans. (A, B, C)**

**Sol.** According to given data  
 $F(x) < 0 \quad \forall x \in (1, 3)$

$$f(x) = x F(x)$$

$$f'(x) = F(x) + x F'(x) \quad \dots(i)$$

$$f'(1) = F(1) + F'(1) < 0$$

(use  $F(1) = 0$  &  $F'(x) < 0$ )

$$f(2) = 2 F(2) < 0$$

(use  $F(x) < 0 \quad \forall x \in (1, 3)$ )

$$f'(x) = F(x) + x F'(x) < 0$$

(use  $F(x) < 0 \quad \forall x \in (1, 3)$ )

$$F'(x) < 0$$

**32. Ans. (C, D)**

**Sol.** Given

$$\int_1^3 x^2 F'(x) dx = -12$$

$$\Rightarrow [x^2 F(x)]_1^3 - 2 \int_1^3 x F(x) dx = -12$$

$$\Rightarrow \int_1^3 f(x) dx = -12 \quad \text{Use } x F(x) = f(x)$$

Given

$$\int_1^3 x^3 F''(x) dx = 40$$

$$\Rightarrow [x^3 F'(x)]_1^3 - 3 \int_1^3 x^2 F'(x) dx = 40$$

$$\Rightarrow [x^2 (f'(x) - F(x))]_1^3 = 4$$

$$9(f(3) - F(3)) - (f'(1) - F(1)) = 4$$

$$9f(3) + 36 - f'(1) = 4$$

$$9f(3) - f'(1) + 32 = 0$$

**33. Ans. (A, C)**

**Sol.** Given that  $f : [a, b] \rightarrow [1, \infty)$

$$g(x) = \begin{cases} 0 & x < a \\ \int_a^x f(t) dt, & a \leq x \leq b \\ \int_a^b f(t) dt, & x > b \end{cases}$$

Now  $g(a^-) = 0 = g(a^+) = g(a)$

$$g(b^-) = g(b^+) = g(b) = \int_a^b f(t) dt$$

$\Rightarrow$   $g$  is continuous  $\forall x \in \mathbb{R}$

$$\text{Now } g'(x) = \begin{cases} 0 & : x < a \\ f(x) & : a < x < b \\ 0 & : x > b \end{cases}$$

$g'(a^-) = 0$  but  $g'(a^+) = f(a) \geq 1$

$\Rightarrow$   $g$  is non differentiable at  $x = a$

and  $g'(b^+) = 0$  but  $g'(b^-) = f(b) \geq 1$

$\Rightarrow$   $g$  is non differentiable at  $x = b$

**34. Ans. (A, C, D)**

$$\text{Sol. } f(x) = \int_{\frac{1}{x}}^x \frac{e^{-\left(\frac{t+1}{t}\right)}}{t} dt$$

$$f'(x) = 1 \cdot \frac{e^{-\left(\frac{x+1}{x}\right)}}{x} - \left( \frac{-1}{x^2} \right) e^{-\left(\frac{1}{x} + x\right)} \cdot \frac{1}{x}$$

$$= \frac{e^{-\left(\frac{x+1}{x}\right)}}{x} + \frac{e^{-\left(\frac{x+1}{x}\right)}}{x} = \frac{2e^{-\left(\frac{x+1}{x}\right)}}{x}$$

$\therefore f(x)$  is monotonically increasing on  $(0, \infty)$

$\Rightarrow$  A is correct & B is wrong.

$$\text{Now } f(x) + f\left(\frac{1}{x}\right) = \int_{1/x}^x \frac{e^{-\left(\frac{t+1}{t}\right)}}{t} dt + \int_x^{1/x} \frac{e^{-\left(\frac{t+1}{t}\right)}}{t} dt$$

$$= 0 \quad \forall x \in (0, \infty)$$

$$\text{Now let } g(x) = f(2^x) = \int_{2^{-x}}^{2^x} \frac{e^{-\left(\frac{t+1}{t}\right)}}{t} dt$$

$$g(-x) = f(2^{-x}) = \int_{2^x}^{2^{-x}} \frac{e^{-\left(\frac{t+1}{t}\right)}}{t} dt = -g(x)$$

$\therefore f(2^x)$  is an odd function.

**35. Ans. (2)**

**Sol.** using integration by part

$$\begin{aligned} & \int_0^1 4x^3 \left( (1-x^2)^5 \right)' dx \\ &= 4x^3 \left( (1-x^2)^5 \right) \Big|_0^1 - \int_0^1 12x^2 (1-x^2)^5 dx \end{aligned}$$

using integration by part

$$= -12 \left[ x^2 \left( (1-x^2)^5 \right) \right]_0^1 - \int_0^1 2x (1-x^2)^5 dx$$

$$= 12.2 \int_0^1 x (1-x^2)^5 dx$$

$$\text{Let } 1-x^2 = t \Rightarrow x dx = -\frac{dt}{2}$$

$$= 24 \int_1^0 t^5 \left( -\frac{dt}{2} \right)$$

$$= 12 \int_0^1 t^5 dt = 2$$

**36. Ans. (A)**

**Sol.** Let  $\cosec x + \cot x = e^u$

$$\cosec x - \cot x = e^{-u}$$

$$\cosec x = \frac{1}{2} (e^u + e^{-u}) \text{ & } \cot x = \frac{1}{2} (e^u - e^{-u})$$

$$\cosec^2 x dx = -\frac{1}{2} (e^u + e^{-u}) du$$

$$\Rightarrow \int_{\ln(\sqrt{2}+1)}^0 2^{17} \left( \frac{1}{2} (e^u + e^{-u}) \right)^{15} \left\{ -\frac{1}{2} (e^u + e^{-u}) \right\} du$$

$$= \int_0^{\ln(\sqrt{2}+1)} 2(e^u + e^{-u})^{16} du.$$

**37. Ans. (B)**

$$\text{Sol. } f(x) = \int_0^{x^2} f(\sqrt{t}) dt$$

$$f'(x) = 2x f(x) \quad \because x \in [0, 2]$$

$$\Rightarrow f'(x) = 2x f(x) \quad \because F'(x) = f'(x)$$

$$\Rightarrow \frac{f'(x)}{f(x)} = 2x \Rightarrow \ln(f(x)) = x^2 + c$$

$$c = 0 \quad (\because f(0) = 1)$$

$$\Rightarrow f(x) = e^{x^2}$$

$$f(x) = \int_0^{x^2} f(\sqrt{t}) dt = \int_0^{x^2} e^t \cdot dt$$

$$\therefore f(2) = \int_0^{x^2} e^t dt = e^4 - 1.$$

**38. Ans. (A)**

**Sol.**

$$\begin{aligned} g\left(\frac{1}{2}\right) &= \lim_{h \rightarrow 0^+} \int_h^{1-h} \frac{dt}{\sqrt{t(1-t)}} \\ &= \lim_{h \rightarrow 0^+} \int_h^{1-h} \frac{dt}{\sqrt{\frac{1}{4} - (t - \frac{1}{2})^2}} \\ &= \sin^{-1} \left( \frac{t - \frac{1}{2}}{\frac{1}{2}} \right) \Big|_0^1 = \sin^{-1}(1) - \sin^{-1}(-1) = \pi \end{aligned}$$

**39. Ans. (D)**

**Sol.** Given,

$$g(a) = \lim_{h \rightarrow 0^+} \int_h^{1-h} t^{-a} (1-t)^{a-1} dt$$

$$g^1(a) = \lim_{h \rightarrow 0^+} \int_h^{1-h} t^{-a} (1-t)^{a-1} (-\ln t + \ln(1-t)) dt$$

$$g'\left(\frac{1}{2}\right) = \lim_{h \rightarrow 0^+} \int_h^{1-h} \frac{\ell n\left(\frac{1-t}{t}\right) dt}{\sqrt{t(1-t)}} \quad \dots(i)$$

$$g^1\left(\frac{1}{2}\right) \lim_{h \rightarrow 0^+} \int_h^{1-h} \frac{\ln\left(\frac{1-(1-t)}{1-t}\right) dt}{\sqrt{(1-t)t}} \quad \dots(ii)$$

$$(Apply \int_a^b f(x) dx = \int_a^b f(a+b-x) dx)$$

$$\Rightarrow 2g^1\left(\frac{1}{2}\right) = \lim_{h \rightarrow 0^+} \int_h^{1-h} 0 dt \Rightarrow g^1\left(\frac{1}{2}\right) = 0$$

**40. Ans. (D)**

**Sol.** (P) let  $f(x) = ax^2 + bx + c$  where  
a,b,c are non negative integers

$$f(0) = c = 0 \quad \dots(1)$$

$$\text{and } \int_0^1 (ax^2 + bx) dx = 1$$

$$= \left[ \frac{ax^3}{3} + \frac{bx^2}{2} \right]_0^1 = \frac{a}{3} + \frac{b}{2} = 1$$

$$\Rightarrow 2a + 3b = 6$$

$$\Rightarrow a = 3 \text{ & } b = 0 \text{ OR } a = 0 \text{ & } b = 2$$

(Q) maximum of  $\sin x^2 + \cos x^2 = \sqrt{2}$

$$\Rightarrow \sin\left(\frac{\pi}{4} + x^2\right) = 1 \text{ but } x^2 \in [0, 13]$$

$$\Rightarrow \frac{\pi}{4} + x^2 = (4n+1)\frac{\pi}{2}$$

$\Rightarrow$  which is satisfied for  $n = 0$  & 1

$\Rightarrow$  4 solutions

$$(R) I = \int_{-2}^2 \frac{3x^2}{1+e^x} dx \text{ put } x$$

$$= -t \quad I = - \int_2^{-2} \frac{3t^2 e^t dt}{1+e^t}$$

$$\Rightarrow 2I = \int_{-2}^2 \frac{3x^2 (1+e^x)}{1+e^x} dx = 2 \int_0^2 3x^2 dx$$

$$\Rightarrow I = \left[ x^3 \right]_0^2 = 8$$

$$(S) \quad \frac{\int_{-1/2}^{1/2} \cos 2x \log\left(\frac{1+x}{1-x}\right) dx}{\int_0^{1/2} \cos 2x \log\left(\frac{1+x}{1-x}\right) dx}$$

$$\begin{aligned} &= \frac{\int_{-1/2}^{1/2} (\text{odd function}) dx}{\int_0^{1/2} \cos 2x \log\left(\frac{1+x}{1-x}\right) dx} = 0 \end{aligned}$$