

PARABOLA

- Let P be a point on the parabola  $y^2 = 4ax$ , where  $a > 0$ . The normal to the parabola at P meets the x-axis at a point Q. The area of the triangle PFQ, where F is the focus of the parabola, is 120. If the slope m of the normal and a are both positive integers, then the pair (a,m) is **[JEE(Advanced) 2023]**  
 (A) (2, 3)                      (B) (1, 3)                      (C) (2, 4)                      (D) (3, 4)
- Consider the parabola  $y^2 = 4x$ . Let S be the focus of the parabola. A pair of tangents drawn to the parabola from the point  $P = (-2, 1)$  meet the parabola at  $P_1$  and  $P_2$ . Let  $Q_1$  and  $Q_2$  be points on the lines  $SP_1$  and  $SP_2$  respectively such that  $PQ_1$  is perpendicular to  $SP_1$  and  $PQ_2$  is perpendicular to  $SP_2$ . Then, which of the following is/are TRUE? **[JEE(Advanced) 2022]**  
 (A)  $SQ_1 = 2$                       (B)  $Q_1Q_2 = \frac{3\sqrt{10}}{5}$                       (C)  $PQ_1 = 3$                       (D)  $SQ_2 = 1$
- Let E denote the parabola  $y^2 = 8x$ . Let  $P = (-2, 4)$ , and let Q and Q' be two distinct points on E such that the lines PQ and PQ' are tangents to E. Let F be the focus of E. Then which of the following statements is (are) TRUE? **[JEE(Advanced) 2021]**  
 (A) The triangle PFQ is a right-angled triangle  
 (B) The triangle QPQ' is a right-angled triangle  
 (C) The distance between P and F is  $5\sqrt{2}$   
 (D) F lies on the line joining Q and Q'

Question Stem for Questions Nos. 4 and 5

Question Stem

Consider the region  $R = \{(x, y) \in \mathbb{R} \times \mathbb{R} : x \geq 0 \text{ and } y^2 \leq 4 - x\}$ . Let F be the family of all circles that are contained in R and have centers on the x-axis. Let C be the circle that has largest radius among the circles in F. Let  $(\alpha, \beta)$  be a point where the circle C meets the curve  $y^2 = 4 - x$ .

- The radius of the circle C is \_\_\_\_\_. **[JEE(Advanced) 2021]**
- The value of  $\alpha$  is \_\_\_\_\_. **[JEE(Advanced) 2021]**
- If a chord, which is not a tangent, of the parabola  $y^2 = 16x$  has the equation  $2x + y = p$ , and midpoint (h, k), then which of the following is(are) possible value(s) of p, h and k? **[JEE(Advanced) 2017]**  
 (A)  $p = 5, h = 4, k = -3$                       (B)  $p = -1, h = 1, k = -3$   
 (C)  $p = -2, h = 2, k = -4$                       (D)  $p = 2, h = 3, k = -4$
- Let P be the point on the parabola  $y^2 = 4x$  which is at the shortest distance from the center S of the circle  $x^2 + y^2 - 4x - 16y + 64 = 0$ . Let Q be the point on the circle dividing the line segment SP internally. Then-  
 (A)  $SP = 2\sqrt{5}$   
 (B)  $SQ : QP = (\sqrt{5} + 1) : 2$   
 (C) the x-intercept of the normal to the parabola at P is 6  
 (D) the slope of the tangent to the circle at Q is  $\frac{1}{2}$  **[JEE(Advanced) 2016]**

8. If the normals of the parabola  $y^2 = 4x$  drawn at the end points of its latus rectum are tangents to the circle  $(x - 3)^2 + (y + 2)^2 = r^2$ , then the value of  $r^2$  is **[JEE(Advanced) 2015]**
9. Let the curve C be the mirror image of the parabola  $y^2 = 4x$  with respect to the line  $x + y + 4 = 0$ . If A and B are the points of intersection of C with the line  $y = -5$ , then the distance between A and B is **[JEE(Advanced) 2015]**
10. Let P and Q be distinct points on the parabola  $y^2 = 2x$  such that a circle with PQ as diameter passes through the vertex O of the parabola. If P lies in the first quadrant and the area of the triangle  $\Delta OPQ$  is  $3\sqrt{2}$ , then which of the following is(are) the coordinates of P ? **[JEE(Advanced) 2015]**
- (A)  $(4, 2\sqrt{2})$                       (B)  $(9, 3\sqrt{2})$                       (C)  $\left(\frac{1}{4}, \frac{1}{\sqrt{2}}\right)$                       (D)  $(1, \sqrt{2})$
11. The common tangents to the circle  $x^2 + y^2 = 2$  and the parabola  $y^2 = 8x$  touch the circle at the point P, Q and the parabola at the points R, S. Then the area of the quadrilateral PQRS is - **[JEE(Advanced) 2014]**
- (A) 3                                      (B) 6                                      (C) 9                                      (D) 15

**Paragraph For Questions 12 and 13**

Let  $a, r, s, t$  be nonzero real numbers. Let  $P(at^2, 2at)$ ,  $Q, R(ar^2, 2ar)$  and  $S(as^2, 2as)$  be distinct points on the parabola  $y^2 = 4ax$ . Suppose that PQ is the focal chord and lines QR and PK are parallel, where K is the point  $(2a, 0)$ .

12. The value of  $r$  is- **[JEE(Advanced) 2014]**
- (A)  $-\frac{1}{t}$                                       (B)  $\frac{t^2 + 1}{t}$                                       (C)  $\frac{1}{t}$                                       (D)  $\frac{t^2 - 1}{t}$
13. If  $st = 1$ , then the tangent at P and the normal at S to the parabola meet at a point whose ordinate is- **[JEE(Advanced) 2014]**
- (A)  $\frac{(t^2 + 1)^2}{2t^3}$                                       (B)  $\frac{a(t^2 + 1)^2}{2t^3}$                                       (C)  $\frac{a(t^2 + 1)^2}{t^3}$                                       (D)  $\frac{a(t^2 + 2)^2}{t^3}$

SOLUTIONS


1. Ans. (A)

Sol. Let point P (at<sup>2</sup>, 2at)

normal at P is y = -tx + 2at + at<sup>3</sup>

y = 0, x = 2a + at<sup>2</sup>

Q(2a + at<sup>2</sup>, 0)



$$\text{Area of } \Delta PFQ = \left| \frac{1}{2} (a + at^2)(2at) \right| = 120$$


$$\therefore m = -t$$

$$\therefore a^2 [1 + m^2] m = 120$$

(a, m) = (2, 3) will satisfy

2. Ans. (B, C, D)

Sol. Let equation of tangent with slope 'm' be



$$T : y = mx + \frac{1}{m}$$

T : passes through (-2, 1) so

$$1 = -2m + \frac{1}{m}$$

$$\Rightarrow m = -1 \text{ or } m = \frac{1}{2}$$

Points are given by  $\left( \frac{a}{m^2}, \frac{2a}{m} \right)$

So, one point will be (1, -2) & (4, 4)

Let P<sub>1</sub>(4, 4) & P<sub>2</sub>(1, -2)

P<sub>1</sub>S : 4x - 3y - 4 = 0

P<sub>2</sub>S : x - 1 = 0

$$PQ_1 = \left| \frac{4(-2) - 3(1) - 4}{5} \right| = 3$$


SP =  $\sqrt{10}$  ; PQ<sub>2</sub> = 3 ; SQ<sub>1</sub> = 1 = SQ<sub>2</sub>

$$\frac{1}{2} \left( \frac{Q_1 Q_2}{2} \right) \times \sqrt{10} = \frac{1}{2} \times 3 \times 1$$

(comparing Areas)

$$\Rightarrow Q_1 Q_2 = \frac{2 \times 3}{\sqrt{10}} = \frac{3\sqrt{10}}{5}$$

3. Ans. (A, B, D)



Sol.


Note that P lies on directrix so triangle PQQ' is right angled, hence QQ' passes through focus F.

$$PF = 4\sqrt{2}$$

Equation of QF is y = x - 2 & PF is x + y = 2

Hence A, B, D.

4. Ans. (1.50)



Sol.

Let the circle be

$$x^2 + y^2 + \lambda x = 0$$

For point of intersection of circle & parabola

$$y^2 = 4 - x.$$

$$x^2 + 4 - x + \lambda x = 0 \Rightarrow x^2 + x(\lambda - 1) + 4 = 0$$

$$\text{For tangency : } \Delta = 0 \Rightarrow (\lambda - 1)^2 - 16 = 0$$

$$\Rightarrow \lambda = 5 \text{ (rejected) or } \lambda = -3$$

$$\text{Circle : } x^2 + y^2 - 3x = 0$$

$$\text{Radius} = \frac{3}{2} = 1.5$$

5. Ans. (2.00)

Sol. For point of intersection :  
 $x^2 - 4x + 4 = 0 \Rightarrow x = 2$  so  $\alpha = 2$

6. Ans. (D)

Sol. Equation of chord with mid point (h, k) :

$$k \cdot y - 16 \left( \frac{x+h}{2} \right) = k^2 - 16h$$


$$\Rightarrow 8x - ky + k^2 - 8h = 0$$

Comparing with  $2x + y - p = 0$ , we get

$$k = -4; 2h - p = 4$$

only (D) satisfies above relation.

7. Ans. (A, C, D)



Sol.

$$y^2 = 4x$$

point P lies on normal to parabola passing through centre of circle

$$y + tx = 2t + t^3 \quad \dots(i)$$

$$8 + 2t = 2t + t^3$$

$$t = 2$$

$$P(4, 4)$$

$$SP = \sqrt{(4-2)^2 + (4-8)^2}$$

$$SP = 2\sqrt{5}$$

$$SQ = 2$$

$$\Rightarrow PQ = 2\sqrt{5} - 2$$

$$\frac{SQ}{QP} = \frac{1}{\sqrt{5}-1} = \frac{\sqrt{5}+1}{4}$$

To find x intercept

put  $y = 0$  in (i)


$$\Rightarrow x = 2 + t^2$$

$$x = 6$$

$\therefore$  Slope of common normal  $= -t = -2$

$\therefore$  Slope of tangent  $= \frac{1}{2}$

8. Ans. (2)



Sol.

The co-ordinates of latus rectum are (1,2) and (1,-2)

clearly slope of tangent is given by  $\frac{dy}{dx} = \frac{2}{y}$

$\therefore$  At  $y = 2$  slope of normal  $= -1$

and At  $y = -2$  slope of normal  $= 1$

$\therefore$  Equation of normal at (1,2)

$$(y - 2) = -1(x - 1) \Rightarrow x + y = 3$$

Now, this line is tangent to circle

$$(x - 3)^2 + (y + 2)^2 = r^2$$

$\therefore$  perpendicular distance from centre to line  
 $=$  Radius of circle

$$\therefore \frac{|3 - 2 - 3|}{\sqrt{2}} = r \Rightarrow r^2 = 2$$

9. Ans. (4)

Sol. Let there be a point  $(t^2, 2t)$  on  $y^2 = 4x$

Clearly its reflection in  $x + y + 4 = 0$  is given by

$$\frac{x - t^2}{1} = \frac{y - 2t}{1} = \frac{-2(t^2 + 2t + 4)}{2}$$


$$\therefore x = -(2t + 4) \text{ \& } y = -(t^2 + 4)$$

$$\text{Now, } y = -5 \Rightarrow t = \pm 1$$

$$\therefore x = -6 \text{ or } x = -2$$

$\therefore$  Distance between A & B = 4

10. Ans. (A, D)



Sol.

$$\because \angle POQ = \frac{\pi}{2} \quad \Rightarrow \quad t_1 t_2 = -4$$

$$\therefore \begin{vmatrix} 0 & 0 & 1 \\ \frac{1}{2} \frac{t_1^2}{2} & t_1 & 1 \\ \frac{1}{2} \frac{t_2^2}{2} & t_2 & 1 \end{vmatrix} = 3\sqrt{2}$$

$$\Rightarrow \left| \frac{t_1^2 t_2 - t_1 t_2^2}{2} \right| = 6\sqrt{2}$$

$$\Rightarrow |t_1 - t_2| = 3\sqrt{2}$$


$$\Rightarrow t_1 + \frac{4}{t_1} = 3\sqrt{2} \quad (\because t_1 > 0)$$

We get  $t_1 = 2\sqrt{2}, \sqrt{2}$

$P(4, 2\sqrt{2})$  or  $(1, \sqrt{2})$

11. Ans. (D)

Sol.



$$y = mx + \frac{2}{m}$$

$$\left| \frac{0 - 0 + \frac{2}{m}}{\sqrt{1 + m^2}} \right| = \sqrt{2} \Rightarrow 2 = m^2(1 + m^2)$$

$$\Rightarrow m = \pm 1$$

$$TP : -x + y = 2$$

So  $P(-1, 1)$  &  $Q(-1, -1)$

&  $R\left(\frac{2}{m}, \frac{4}{m}\right) \equiv R(2, 4)$  &  $S(2, -4)$

$$\text{So } \Delta = \frac{1}{2} \cdot 10.3 = 15$$


So  $P(-1, 1)$  &  $Q(-1, -1)$

&  $R\left(\frac{2}{m}, \frac{4}{m}\right) \equiv R(2, 4)$  &  $S(2, -4)$

$$\text{So } \Delta = \frac{1}{2} \cdot 10.3 = 15$$

12. Ans. (D)

Sol.



$\because$  PQ is a focal chord

$\therefore$  co-ordinates of point Q are  $\left(\frac{a}{t^2}, -\frac{2a}{t}\right)$

$$m_{QR} = \frac{2a\left(r + \frac{1}{t}\right)}{a\left(r^2 - \frac{1}{t^2}\right)} = \frac{2}{\left(r - \frac{1}{t}\right)}$$

$$m_{PK} = \frac{2at - 0}{a(t^2 - 2)} = \frac{2t}{t^2 - 2}$$

Given  $m_{QR} = m_{PK}$

$$\Rightarrow \frac{2}{r - \frac{1}{t}} = \frac{2t}{t^2 - 2} \Rightarrow r = \frac{t^2 - 2}{t} + \frac{1}{t}$$

$$\Rightarrow r = t - \frac{2}{t} + \frac{1}{t} \Rightarrow r = \frac{t^2 - 1}{t}$$

13. Ans. (B)

Sol. Equation of tangent at point P is

$$ty = x + at^2 \quad \dots(i)$$

Equation of normal at point S is

$$\frac{1}{t}x + y = \frac{2a}{t} + \frac{a}{t^3}$$

$$\Rightarrow t^2x + t^3y = 2at^2 + a \quad \dots(ii)$$

Multiply equation (i) by  $t^2$  and then subtract from equation (ii),

we get,

$$2t^3y = 2at^2 + at^4 + a$$

$$\Rightarrow 2t^3y = a(1 + t^4 + 2t^2)$$

$$\Rightarrow y = \frac{a(1 + t^2)^2}{2t^3}$$