

SEQUENCE & SERIES

1. Let $\overbrace{75\dots57}^r$ denote the $(r + 2)$ digit number where the first and the last digits are 7 and the remaining r digits are 5. Consider the sum $S = 77 + 757 + 7557 + \dots + \overbrace{75\dots57}^{98}$. If $S = \frac{\overbrace{75\dots57}^{99} + m}{n}$, where m and n are natural numbers less than 3000, then the value of $m + n$ is **[JEE(Advanced) 2023]**
2. Let l_1, l_2, \dots, l_{100} be consecutive terms of an arithmetic progression with common difference d_1 , and let w_1, w_2, \dots, w_{100} be consecutive terms of another arithmetic progression with common difference d_2 , where $d_1 d_2 = 10$. For each $i = 1, 2, \dots, 100$, let R_i be a rectangle with length l_i , width w_i and area A_i . If $A_{51} - A_{50} = 1000$, then the value of $A_{100} - A_{90}$ is **[JEE(Advanced) 2022]**
3. Let a_1, a_2, a_3, \dots be an arithmetic progression with $a_1 = 7$ and common difference 8. Let T_1, T_2, T_3, \dots be such that $T_1 = 3$ and $T_{n+1} - T_n = a_n$ for $n \geq 1$. Then, which of the following is/are TRUE ? **[JEE(Advanced) 2022]**

(A) $T_{20} = 1604$

(B) $\sum_{k=1}^{20} T_k = 10510$

(C) $T_{30} = 3454$

(D) $\sum_{k=1}^{30} T_k = 35610$

4. Let m be the minimum possible value of $\log_3(3^{y_1} + 3^{y_2} + 3^{y_3})$, where y_1, y_2, y_3 are real numbers for which $y_1 + y_2 + y_3 = 9$. Let M be the maximum possible value of $(\log_3 x_1 + \log_3 x_2 + \log_3 x_3)$, where x_1, x_2, x_3 are positive real numbers for which $x_1 + x_2 + x_3 = 9$. Then the value of $\log_2(m^3) + \log_3(M^2)$ is **[JEE(Advanced) 2020]**

5. Let a_1, a_2, a_3, \dots be a sequence of positive integers in arithmetic progression with common difference 2. Also, let b_1, b_2, b_3, \dots be a sequence of positive integers in geometric progression with common ratio 2. If $a_1 = b_1 = c$, then the number of all possible values of c , for which the equality

$$2(a_1 + a_2 + \dots + a_n) = b_1 + b_2 + \dots + b_n$$

holds for some positive integer n , is **[JEE(Advanced) 2020]**

6. Let α and β be the roots of $x^2 - x - 1 = 0$, with $\alpha > \beta$. For all positive integers n , define

$$a_n = \frac{\alpha^n - \beta^n}{\alpha - \beta}, \quad n \geq 1$$

$$b_1 = 1 \text{ and } b_n = a_{n-1} + a_{n+1}, \quad n \geq 2.$$

Then which of the following options is/are correct ? **[JEE(Advanced) 2019]**

(A) $a_1 + a_2 + a_3 + \dots + a_n = a_{n+2} - 1$ for all $n \geq 1$ (B) $\sum_{n=1}^{\infty} \frac{a_n}{10^n} = \frac{10}{89}$

(C) $\sum_{n=1}^{\infty} \frac{b_n}{10^n} = \frac{8}{89}$ (D) $b_n = \alpha^n + \beta^n$ for all $n \geq 1$

7. Let AP (a; d) denote the set of all the terms of an infinite arithmetic progression with first term a and common difference $d > 0$. If $AP(1; 3) \cap AP(2; 5) \cap AP(3; 7) = AP(a; d)$ then $a + d$ equals _____.
[JEE(Advanced) 2019]
8. Let X be the set consisting of the first 2018 terms of the arithmetic progression 1, 6, 11,, and Y be the set consisting of the first 2018 terms of the arithmetic progression 9, 16, 23,, Then, the number of elements in the set $X \cup Y$ is _____.
[JEE(Advanced) 2018]
9. The sides of the right angled triangle are in arithmetic progression. If the triangle has area 24, then what is the length of its smallest side ?
[JEE(Advanced) 2017]
10. Let $b_i > 1$ for $i = 1, 2, \dots, 101$. Suppose $\log_e b_1, \log_e b_2, \dots, \log_e b_{101}$ are in Arithmetic Progression (A.P.) with the common difference $\log_e 2$. Suppose a_1, a_2, \dots, a_{101} are in A.P. such that $a_1 = b_1$ and $a_{51} = b_{51}$. If $t = b_1 + b_2 + \dots + b_{51}$ and $s = a_1 + a_2 + \dots + a_{51}$ then
[JEE(Advanced) 2016]
 (A) $s > t$ and $a_{101} > b_{101}$ (B) $s > t$ and $a_{101} < b_{101}$
 (C) $s < t$ and $a_{101} > b_{101}$ (D) $s < t$ and $a_{101} < b_{101}$
11. Suppose that all the terms of an arithmetic progression (A.P.) are natural numbers. If the ratio of the sum of the first seven terms to the sum of the first eleven terms is $6 : 11$ and the seventh term lies in between 130 and 140, then the common difference of this A.P. is
[JEE(Advanced) 2015]
12. Let a,b,c be positive integers such that $\frac{b}{a}$ is an integer. If a,b,c are in geometric progression and the arithmetic mean of a,b,c is $b + 2$, then the value of $\frac{a^2 + a - 14}{a + 1}$ is
[JEE(Advanced) 2014]

SOLUTIONS

1. **Ans. (1219)**

Sol. $S = 77 + 757 + 7557 + \dots + \underbrace{75\dots57}_{98}$

$10S = 770 + 7570 + \dots + 75 \dots 570 + 755 \dots 570$

$$9S = -77 + \underbrace{13+13+\dots+13}_{98 \text{ times}} + \underbrace{75\dots570}_{98}$$

$$= -77 + 13 \times 98 + \underbrace{75\dots57}_{99} + 13$$

$$S = \frac{\underbrace{75\dots57}_{99} + 1210}{9}$$

$m = 1210$

$n = 9$

$m + n = 1219$

2. **Ans. (18900.00)**

Sol. Given

$$A_{51} - A_{50} = 1000 \Rightarrow \ell_{51}w_{51} - \ell_{50}w_{50} = 1000$$

$$\Rightarrow (\ell_1 + 50d_1)(w_1 + 50d_2) - (\ell_1 + 49d_1)(w_1 + 49d_2) = 1000$$

$$\Rightarrow (\ell_1 d_2 + w_1 d_1) = 10 \quad \dots(1)$$

(As $d_1 d_2 = 10$)

$$\therefore A_{100} - A_{90} = \ell_{100}w_{100} - \ell_{90}w_{90}$$

$$= (\ell_1 + 99d_1)(w_1 + 99d_2) - (\ell_1 + 89d_1)(w_1 + 89d_2)$$

$$= 10(\ell_1 d_2 + w_1 d_1) + (99^2 - 89^2)d_1 d_2$$

$$= 10(10) + \underbrace{(99-89)}_{=10}(99+89)(10)$$

(As, $d_1 d_2 = 10$)

$$= 100(1 + 188) = 100(189) = 18900$$

3. **Ans. (B, C)**

Sol. $a_1 = 7, d = 8$

$$T_{n+1} - T_n = a_n \forall n \geq 1$$

$$S_n = T_1 + T_2 + T_3 + \dots + T_{n-1} + T_n$$

$$S_n = T_1 + T_2 + T_3 + \dots + T_{n-1} + T_n$$

on subtraction

$$T_n = T_1 + a_1 + a_2 + \dots + a_{n-1}$$

$$T_n = 3 + (n-1)(4n-1)$$

$$T_n = 4n^2 - 5n + 4$$

$$\sum_{k=1}^n T_k = 4 \sum_{k=1}^n k^2 - 5 \sum_{k=1}^n k + 4n$$

$$T_{20} = 1504$$

$$T_{30} = 3454$$

$$\sum_{k=1}^{30} T_k = 35615$$

$$\sum_{k=1}^{20} T_k = 10510$$

4. **Ans. (8.00)**

Sol. $\frac{3^{y_1} + 3^{y_2} + 3^{y_3}}{3} \geq \left[3^{(y_1+y_2+y_3)} \right]^{\frac{1}{3}}$

$$\Rightarrow 3^{y_1} + 3^{y_2} + 3^{y_3} \geq 3^4$$

$$\Rightarrow \log_3(3^{y_1} + 3^{y_2} + 3^{y_3}) \geq 41$$

$$\Rightarrow m = 4$$

Also, $\frac{x_1 + x_2 + x_3}{3} \geq \sqrt[3]{x_1 x_2 x_3}$

$$\Rightarrow x_1 x_2 x_3 \leq 27$$

$$\Rightarrow \log_3 x_1 + \log_3 x_2 + \log_3 x_3 \leq 3$$

$$\Rightarrow M = 3$$

Thus, $\log_2(m^3) + \log_3(M^2) = 6 + 2 = 8$

5. **Ans. (1.00)**

Sol. Given $2(a_1 + a_2 + \dots + a_n) = b_1 + b_2 + \dots + b_n$

$$\Rightarrow 2 \times \frac{n}{2}(2c + (n-2)c) = c \left(\frac{2^n - 1}{2 - 1} \right)$$

$$\Rightarrow 2n^2 - 2n = c(2^n - 1 - 2n)$$

$$\Rightarrow c = \frac{2n^2 - 2n}{2^n - 1 - 2n} \in \mathbb{N}$$

So, $2n^2 - 2n \geq 2^n - 1 - 2n$

$$\Rightarrow 2n^2 + 1 \geq 2^n \Rightarrow n < 7$$

$$\Rightarrow n \text{ can be } 1, 2, 3, \dots$$

Checking c against these values of n

we get $c = 12$ (when $n = 3$)

Hence number of such $c = 1$

6. **Ans. (A, B, D)**

Sol. α, β are roots of $x^2 - x - 1$

$$a_{r+2} - a_r = \frac{(\alpha^{r+2} - \beta^{r+2}) - (\alpha^r - \beta^r)}{\alpha - \beta}$$

$$= \frac{(\alpha^{r+2} - \alpha^r) - (\beta^{r+2} - \beta^r)}{\alpha - \beta}$$

$$= \frac{\alpha^r(\alpha^2 - 1) - \beta^r(\beta^2 - 1)}{\alpha - \beta}$$

$$= \frac{\alpha^r \alpha - \beta^r \beta}{\alpha - \beta} = \frac{\alpha^{r+1} - \beta^{r+1}}{\alpha - \beta} = a_{r+1}$$

$$\Rightarrow a_{r+2} - a_{r+1} = a_r$$

$$\Rightarrow \sum_{r=1}^n a_r = a_{n+2} - a_2 = a_{n+2} - \frac{\alpha^2 - \beta^2}{\alpha - \beta}$$

$$= a_{n+2} - (\alpha + \beta) = a_{n+2} - 1$$

Now $\sum_{n=1}^{\infty} \frac{a_n}{10^n} = \frac{\sum_{n=1}^{\infty} \left(\frac{\alpha}{10}\right)^n - \sum_{n=1}^{\infty} \left(\frac{\beta}{10}\right)^n}{\alpha - \beta}$

$$\frac{\frac{\alpha}{10} - \frac{\beta}{10}}{1 - \frac{\alpha}{10} - 1 + \frac{\beta}{10}} = \frac{\frac{\alpha}{10 - \alpha} - \frac{\beta}{10 - \beta}}{\alpha - \beta}$$

$$= \frac{10}{(10 - \alpha)(10 - \beta)} = \frac{10}{89}$$

$$\sum_{n=1}^{\infty} \frac{b_n}{10^n} = \sum_{n=1}^{\infty} \frac{a_{n-1} + a_{n+1}}{10^n} = \frac{\frac{\alpha}{10} + \frac{\beta}{10}}{1 - \frac{\alpha}{10} - 1 + \frac{\beta}{10}} = \frac{12}{89}$$

Further, $b_n = a_{n-1} + a_{n+1}$

$$= \frac{(\alpha^{n-1} - \beta^{n-1}) + (\alpha^{n+1} - \beta^{n+1})}{\alpha - \beta}$$

(as $\alpha\beta = -1 \Rightarrow \alpha^{n-1} = -\alpha^n\beta$ & $\beta^{n-1} = -\alpha\beta^n$)

$$= \frac{\alpha^n(\alpha - \beta) + (\alpha - \beta)\beta^n}{\alpha - \beta} = \alpha^n + \beta^n$$

7. Ans. (157.00)

Sol. We equate the general terms of three respective A.P.'s as $1 + 3a = 2 + 5b = 3 + 7c$
 $\Rightarrow 3$ divides $1 + 2b$ and 5 divides $1 + 2c$
 $\Rightarrow 1 + 2c = 5, 15, 25$ etc.
 So, first such terms are possible when $1 + 2c = 15$ i.e. $c = 7$
 Hence, first term = $a = 52$
 $d = \text{lcm}(3, 5, 7) = 105 \Rightarrow a + d = 157$

8. Ans. (3748)

Sol. X : 1, 6, 11,, 10086
 Y : 9, 16, 23,, 14128
 $X \cap Y$: 16, 51, 86,

Let $m = n(X \cap Y)$

$$\therefore 16 + (m - 1) \times 35 \leq 10086$$


$$\Rightarrow m \leq 288.71$$

$$\Rightarrow m = 288$$

$$\therefore n(X \cup Y) = n(X) + n(Y) - n(X \cap Y)$$

$$= 2018 + 2018 - 288 = 3748$$

9. Ans. (6)



where $d > 0, a > 0$
 \Rightarrow length of smallest side = $a - d$
 Now $(a + d)^2 = a^2 + (a - d)^2$
 $\Rightarrow a(a - 4d) = 0$

$$\therefore a = 4d \quad \dots(1)$$

(As $a = 0$ is rejected)

Also, $\frac{1}{2}a(a - d) = 24$

$$\Rightarrow a(a - d) = 48 \quad \dots(2)$$

\therefore From (1) and (2), we get $a = 8, d = 2$

Hence, length of smallest side

$$\Rightarrow (a - d) = (8 - 2) = 6$$

10. Ans. (B)

Sol. If $\log_e b_1, \log_e b_2, \dots, \log_e b_{101} \rightarrow AP$; $D = \log_e 2$

$$\Rightarrow b_1 b_2 b_3 \dots b_{101} \rightarrow GP ; \quad r = 2$$

$$\therefore b_1, 2b_1, 2^2 b_1, \dots, 2^{100} b_1 \dots GP$$

$$a_1 a_2 a_3 \dots a_{101} \dots AP$$

Given, $a_1 = b_1$ & $a_{51} = b_{51}$

$$\Rightarrow a_1 + 50D = 2^{50} b_1$$

$$\therefore a_1 + 50D = 2^{50} a_1 \quad (\text{As } b_1 = a_1)$$

Now, $t = b_1(2^{51} - 1)$; $s = \frac{51}{2}(2a_1 + 50D)$

$$\Rightarrow t < a_1 \cdot 2^{51} \dots(i) \quad ; \quad s = \frac{51}{2}(a_1 + a_1 + 50D)$$

$$s = \frac{51}{2}(a_1 + 2^{50} a_1)$$

$$s = \frac{51a_1}{2} + \frac{51}{2} \cdot 2^{50} a_1$$

$$\Rightarrow s > a_1 \cdot 2^{51} \quad \dots(ii)$$

clearly $s > t$ (from equation (i) and (ii))

Also $a_{101} = a_1 + 100D$; $b_{101} = b_1 \cdot 2^{100}$

$$\therefore a_{101} = a_1 + 100 \left(\frac{2^{50} a_1 - a_1}{50} \right); b_{101}$$

$$= 2^{100} a_1 \quad \dots(\text{iii})$$

$$a_{101} = a_1 + 2^{51} a_1 - 2a_1 \Rightarrow a_{101} = 2^{51} a_1 - a_1$$

$$\Rightarrow a_{101} < 2^{51} a_1 \quad \dots(\text{iv})$$

clearly $b_{101} > a_{101}$ (from equation (iii) and (iv))

11. **Ans. (9)**

$$\text{Sol. } \frac{\frac{7}{2}[2a+6d]}{\frac{11}{2}[2a+10d]} = \frac{6}{11} \Rightarrow \frac{7(2a+6d)}{(2a+10d)} = 6$$

$$\Rightarrow 2a = 18d$$

$$a = 9d$$

$$\text{also } 130 < a + 6d < 140$$

$$\frac{26}{3} < d < \frac{28}{3} \Rightarrow d = 9$$

12. **Ans. (4)**

Sol. Let a, b, c are a, ar, ar^2 where $r \in \mathbb{N}$

$$\text{also } \frac{a+b+c}{3} = b+2$$

$$\Rightarrow a + ar + ar^2 = 3(ar) + 6$$

$$\Rightarrow ar^2 - 2ar + a = 6$$

$$\Rightarrow (r-1)^2 = \frac{6}{a}$$

$\therefore \frac{6}{a}$ must be perfect square & $a \in \mathbb{N}$

$\therefore a$ can be 6 only.

$$\Rightarrow r-1 = \pm 1 \Rightarrow r = 2$$

$$\& \frac{a^2 + a - 14}{a+1} = \frac{36 + 6 - 14}{7} = 4$$