DIFFERENTIABILITY

1. Let S be the set of points where the function, $f(x) = |2 - |x - 3||$, $x \in R$, is not differentiable.

Then $\sum_{x \in S} f(f(x))$ is equal to _____.

2. If a function f(x) defined by

$$f(x) = \begin{cases} ae^x + be^{-x}, & -1 \le x < 1 \\ cx^2, & 1 \le x \le 3 \\ ax^2 + 2cx, & 3 < x \le 4 \end{cases}$$ be continuous

for some a, b, c $\in$ R and $f'(0) + f'(2) = e$, then the value of of a is :

(1) $\frac{e}{e^2 - 3e - 13}$ (2) $\frac{e}{e^2 + 3e + 13}$

(3) $\frac{1}{e^2 - 3e + 13}$ (4) $\frac{e}{e^2 - 3e + 13}$

3. Suppose a differentiable function f(x) satisfies the identity $f(x + y) = f(x) + f(y) + xy^2 + x^2y$, for all real x and y. If $\lim_{x \to 0} \frac{f(x)}{x} = 1$, then $f'(3)$ is equal to _____.

4. The function $f(x) = \begin{cases} \frac{\pi}{4} + \tan^{-1} x, & |x| \le 1 \\ \frac{1}{2}(|x| - 1), & |x| > 1 \end{cases}$ is :

(1) continuous on R–{1} and differentiable on R – {–1, 1}.

(2) both continuous and differentiable on R – {–1}.

(3) continuous on R – {–1} and differentiable on R – {–1, 1}.

(4) both continuous and differentiable on R –{1}

5. If the function $f(x) = \begin{cases} k_1(x - \pi)^2 - 1, & x \le \pi \\ k_2 \cos x, & x > \pi \end{cases}$ is twice differentiable, then the ordered pair (k_1, k_2) is equal to :

(1) $\left(\frac{1}{2}, 1\right)$ (2) (1, 1)

(3) $\left(\frac{1}{2}, -1\right)$ (4) (1, 0)

6. Let $f : R \to R$ be defined as

$$f(x) = \begin{cases} x^5 \sin\left(\frac{1}{x}\right) + 5x^2, & x < 0 \\ 0, & x = 0 \\ x^5 \cos\left(\frac{1}{x}\right) + \lambda x^2, & x > 0 \end{cases}$$. The value

of λ for which $f''(0)$ exists, is _.

7. Let $f : R \to R$ be a function defined by $f(x) = \max\{x, x^2\}$. Let S denote the set of all points in R, where f is not differentiable. Then :

(1) {0, 1} (2) {0}

(3) ϕ(an empty set) (4) {1}

SOLUTION

1. NTA Ans. (3)

Sol. $f(x) = |2 - |x - 3||$

f is not differentiable at

$x = 1, 3, 5$



$\Rightarrow \sum_{x \in s} f(f(x)) = f(f(1)) + f(f(3)) + f(f(5))$

$= f(0) + f(2) + f(0)$

$= 1 + 1 + 1 = 3$

2. Official Ans. by NTA (4)

Sol. $f(x) = \begin{cases} ae^x + be^{-x}, & -1 \le x < 1 \\ cx^2, & 1 \le x \le 3 \\ ax^2 + 2cx, & 3 < x \le 4 \end{cases}$

For continuity at x = 1

$\lim_{x \to 1^-} f(x) = \lim_{x \to 1^+} f(x)$

$\Rightarrow \boxed{ae + be^{-1} = c} \Rightarrow \boxed{b = ce - ae^2}$

...(1)

For continuity at x = 3

$\lim_{x \to 3^-} f(x) = \lim_{x \to 3^+} f(x)$

$\Rightarrow 9c = 9a + 6c$

$\Rightarrow c = 3a$ ...(2)

$f'(0) + f'(2) = e$

$(ae^x - be^x)_{x=0} + (2cx)_{x=2} = e$

$\Rightarrow \boxed{a - b + 4c = e}$ ...(3)

From (1), (2) & (3)

$a - 3ae + ae^2 + 12a = e$

$\Rightarrow a(e^2 + 13 - 3e) = e$

$\Rightarrow a = \dfrac{e}{e^2 - 3e + 13}$

3. Official Ans. by NTA (10)

Sol. Since, $\lim_{x \to 0} \dfrac{f(x)}{x}$ exist $\Rightarrow f(0) = 0$

Now, $f'(x) = \lim_{h \to 0} \dfrac{f(x+h) - f(x)}{h}$

$= \lim_{h \to 0} \dfrac{f(h) + xh^2 + x^2h}{h}$ (take y = h)

$= \lim_{h \to 0} \dfrac{f(h)}{h} + \lim_{h \to 0}(xh) + x^2$

$\Rightarrow f'(x) = 1 + 0 + x^2$

$\Rightarrow f'(3) = 10$

4. Official Ans. by NTA (1)

Sol. $f(x) = \begin{cases} \dfrac{\pi}{4} + \tan^{-1} x, & x \in (-\infty, -1] \cup [1, \infty) \\ -\dfrac{(x+1)}{2}, & x \in (-1, 0] \\ \dfrac{x-1}{2}, & x \in (0, 1) \end{cases}$

for continuity at x = –1

L.H.L. = $\dfrac{\pi}{4} - \dfrac{\pi}{4} = 0$

R.H.L. = 0

so, continuous at x = –1

for continuity at x = 1

L.H.L. = 0

R.H.L. = $\dfrac{\pi}{4} + \dfrac{\pi}{4} = \dfrac{\pi}{2}$

so, not continuous at x = 1

For differentiability at x = –1

L.H.D. = $\dfrac{1}{1+1} = \dfrac{1}{2}$

R.H.D. = $-\dfrac{1}{2}$

so, non differentiable at x = –1

node06\B0BA-BB\Kota\JEE MAIN\Jee Main-2020_Subject Topic PDF With Solution\Mathematics\English\Differentiability.p65

5. Official Ans. by NTA (1)

Sol. f(x) is continuous and differentiable

$f(\pi^-) = f(\pi) = f(\pi^+)$

$-1 = -k_2$

$\boxed{k_2 = 1}$

$$f'(x) = \begin{cases} 2k_1(x-\pi)\ ; x \le \pi \\ -k_2 \sin x \quad ; x > \pi \end{cases}$$

$f'(\pi^-) = f'(\pi^+)$

$0 = 0$

so, differentiable at x = 0

$$f''(x) = \begin{cases} 2k_1 \quad ; x \le \pi \\ -k_2 \cos x\ ; x > \pi \end{cases}$$

$f''(\pi^-) = f''(\pi^+)$

$2k_1 = k_2$

$\boxed{k_1 = \frac{1}{2}}$

$(k_1, k_2) = \left(\frac{1}{2}, 1\right)$

6. Official Ans. by NTA (5.00)

Sol. $f(x) = x^5.\sin\frac{1}{x} + 5x^2$ if $x < 0$

$f(x) = 0$ if $x = 0$

$f(x) = x^5.\cos\frac{1}{x} + \lambda x^2$ if $x > 0$

LHD of $f'(x)$ at x = 0 is 10

RHD of $f'(x)$ at x = 0 is 2λ

if $f''(0)$ exists then

$2\lambda = 10 \Rightarrow \lambda = 5$

7. Official Ans. by NTA (1)

Sol. $f(x) = \max(x, x^2)$



Non-differentiable at x = 0, 1

S = {0, 1}

node06\B0BA-BB\Kota\JEE MAIN\Jee Main-2020_Subject Topic PDF With Solution\Mathamatics\English\Differentiability.p65