

TANGENT & NORMAL

1. The length of the perpendicular from the origin, on the normal to the curve, $x^2 + 2xy - 3y^2 = 0$ at the point (2,2) is
 (1) $4\sqrt{2}$ (2) $2\sqrt{2}$
 (3) 2 (4) $\sqrt{2}$
2. Let the normal at a point P on the curve $y^2 - 3x^2 + y + 10 = 0$ intersect the y-axis at $\left(0, \frac{3}{2}\right)$. If m is the slope of the tangent at P to the curve, then |m| is equal to
3. If the tangent to the curve $y = x + \sin y$ at a point (a, b) is parallel to the line joining $\left(0, \frac{3}{2}\right)$ and $\left(\frac{1}{2}, 2\right)$, then :
 (1) $b = a$ (2) $b = \frac{\pi}{2} + a$
 (3) $|b - a| = 1$ (4) $|a + b| = 1$
4. The equation of the normal to the curve $y = (1+x)^{2y} + \cos^2(\sin^{-1}x)$ at $x = 0$ is :
 (1) $y = 4x + 2$ (2) $x + 4y = 8$
 (3) $y + 4x = 2$ (4) $2y + x = 4$
5. If the surface area of a cube is increasing at a rate of $3.6 \text{ cm}^2/\text{sec}$, retaining its shape; then the rate of change of its volume (in cm^3/sec), when the length of a side of the cube is 10 cm, is :
 (1) 9 (2) 18
 (3) 10 (4) 20
6. If the tangent of the curve, $y = e^x$ at a point (c, e^c) and the normal to the parabola, $y^2 = 4x$ at the point (1, 2) intersect at the same point on the x-axis, then the value of c is _____.
7. If the lines $x + y = a$ and $x - y = b$ touch the curve $y = x^2 - 3x + 2$ at the points where the curve intersects the x-axis, then $\frac{a}{b}$ is equal to _____.
8. The position of a moving car at time t is given by $f(t) = at^2 + bt + c$, $t > 0$, where a, b and c are real numbers greater than 1. Then the average speed of the car over the time interval $[t_1, t_2]$ is attained at the point :
 (1) $a(t_2 - t_1) + b$ (2) $(t_2 - t_1)/2$
 (3) $2a(t_1 + t_2) + b$ (4) $(t_1 + t_2)/2$

SOLUTION

1. NTA Ans. (2)

Sol. $x^2 + 2xy - 3y^2 = 0$

m_N = slope of normal drawn to curve at (2,2)
is -1

$L : x + y = 4.$

perpendicular distance of L from (0,0)

$$= \frac{|0+0-4|}{\sqrt{2}} = 2\sqrt{2}$$

(2) Option

2. NTA Ans. (4.00)

Sol. Let $P(\alpha, \beta)$

so, $\beta^2 - 3\alpha^2 + \beta + 10 = 0$
...(i)

Now, $2yy' - 6x + y' = 0$

$$\Rightarrow m = \frac{6\alpha}{2\beta+1} \dots (ii)$$

Also, $\frac{\beta - \frac{3}{2}}{\alpha} = -\frac{1}{m}$

$$\Rightarrow \frac{2\beta-3}{2\alpha} = -\frac{(2\beta+1)}{6\alpha} \text{ (from (ii))}$$

$$\Rightarrow \beta = 1 \Rightarrow \alpha^2 = 4 \text{ (from (1))}$$

Hence, $|m| = \frac{12}{3} = 4.00$

3. Official Ans. by NTA (3)

Sol. Slope of tangent to the curve $y = x + \sin y$

at (a, b) is $\frac{2 - \frac{3}{2}}{\frac{1}{2} - 0} = 1$

$$\Rightarrow \left. \frac{dy}{dx} \right|_{x=a} = 1$$

$$\frac{dy}{dx} = 1 + \cos y \cdot \frac{dy}{dx} \text{ (from equation of curve)}$$

$$\left. \frac{dy}{dx} \right|_{x=a} = 1 + \cos b \cdot \left. \frac{dy}{dx} \right|_{x=a}$$

$$\Rightarrow \cos b = 0$$

$$\Rightarrow \sin b = \pm 1$$

Now, from curve $y = x + \sin y$

$$b = a + \sin b$$

$$\Rightarrow |b - a| = |\sin b| = 1$$

4. Official Ans. by NTA (2)

Sol. Given equation of curve
 $y = (1+x)^{2y} + \cos^2(\sin^{-1}x)$

at $\boxed{x=0}$

$$y = (1+0)^{2y} + \cos^2(\sin^{-1}0)$$

$$y = 1 + 1$$

$$\boxed{y=2}$$

So we have to find the normal at (0, 2)

Now $y = e^{2y \ln(1+x)} + \cos^2(\cos^{-1} \sqrt{1-x^2})$

$$y = e^{2y \ln(1+x)} + (\sqrt{1-x^2})^2$$

