


LIQUID SOLUTION

- When 9.45 g of ClCH_2COOH is added to 500 mL of water, its freezing point drops by 0.5°C . The dissociation constant of ClCH_2COOH is $x \times 10^{-3}$. The value of x is _____. (Rounded off to the nearest integer)
[$K_{\text{f}(\text{H}_2\text{O})} = 1.86 \text{ K kg mol}^{-1}$]
- C_6H_6 freezes at 5.5°C . The temperature at which a solution 10 g of C_4H_{10} in 200 g of C_6H_6 freeze is _____ $^\circ\text{C}$. (The molal freezing point depression constant of C_6H_6 is 5.12°C/m .)
- 1 molal aqueous solution of an electrolyte A_2B_3 is 60% ionised. The boiling point of the solution at 1 atm is _____ K. (Rounded-off to the nearest integer)
[Given K_{b} for $(\text{H}_2\text{O}) = 0.52 \text{ K kg mol}^{-1}$]
- If a compound AB dissociates to the extent of 75% in an aqueous solution, the molality of the solution which shows a 2.5 K rise in the boiling point of the solution is _____ molal. (Rounded-off to the nearest integer)
[$K_{\text{b}} = 0.52 \text{ K kg mol}^{-1}$]
- 224 mL of $\text{SO}_{2(\text{g})}$ at 298 K and 1 atm is passed through 100 mL of 0.1 M NaOH solution. The non-volatile solute produced is dissolved in 36 g of water. The lowering of vapour pressure of solution (assuming the solution is dilute) ($P_{(\text{H}_2\text{O})} = 24 \text{ mm of Hg}$) is $x \times 10^{-2} \text{ mm of Hg}$, the value of x is _____. (Integer answer)
- When 12.2 g of benzoic acid is dissolved in 100 g of water, the freezing point of solution was found to be -0.93°C ($K_{\text{f}}(\text{H}_2\text{O}) = 1.86 \text{ K kg mol}^{-1}$). The number (n) of benzoic acid molecules associated (assuming 100% association) is _____.
- AB_2 is 10% dissociated in water to A^{2+} and B^- . The boiling point of a 10.0 molal aqueous solution of AB_2 is _____ $^\circ\text{C}$. (Round off to the Nearest Integer).
[Given : Molal elevation constant of water $K_{\text{b}} = 0.5 \text{ K kg mol}^{-1}$ boiling point of pure water $= 100^\circ\text{C}$]
- At 363 K, the vapour pressure of A is 21 kPa and that of B is 18 kPa. One mole of A and 2 moles of B are mixed. Assuming that this solution is ideal, the vapour pressure of the mixture is _____ kPa. (Round of to the Nearest Integer).
- The oxygen dissolved in water exerts a partial pressure of 20 kPa in the vapour above water. The molar solubility of oxygen in water is _____ $\times 10^{-5} \text{ mol dm}^{-3}$. (Round off to the Nearest Integer).
[Given : Henry's law constant $= K_{\text{H}} = 8.0 \times 10^4 \text{ kPa for O}_2$.
Density of water with dissolved oxygen $= 1.0 \text{ kg dm}^{-3}$]
- A 1 molal $\text{K}_4\text{Fe}(\text{CN})_6$ solution has a degree of dissociation of 0.4. Its boiling point is equal to that of another solution which contains 18.1 weight percent of a non electrolytic solute A. The molar mass of A is _____ u. (Round off to the Nearest Integer).
[Density of water $= 1.0 \text{ g cm}^{-3}$]
- 2 molal solution of a weak acid HA has a freezing point of 3.885°C . The degree of dissociation of this acid is _____ $\times 10^{-3}$. (Round off to the Nearest Integer).
[Given : Molal depression constant of water $= 1.85 \text{ K kg mol}^{-1}$ Freezing point of pure water $= 0^\circ\text{C}$]

12. A solute dimerizes in water. The boiling point of a 2 molar solution of A is 100.52°C . The percentage association of A is _____.
(Round off to the Nearest integer)
[Use : K_b for water = $0.52 \text{ K kg mol}^{-1}$
Boiling point of water = 100°C]
13. Which one of the following 0.06 M aqueous solutions has lowest freezing point ?
(1) $\text{Al}_2(\text{SO}_4)_3$ (2) $\text{C}_6\text{H}_{12}\text{O}_6$
(3) KI (4) K_2SO_4
14. CO_2 gas is bubbled through water during a soft drink manufacturing process at 298 K. If CO_2 exerts a partial pressure of 0.835 bar then x mol of CO_2 would dissolve in 0.9 L of water. The value of x is _____.
(Nearest integer)
(Henry's law constant for CO_2 at 298 K is $1.67 \times 10^3 \text{ bar}$)
15. When 3.00 g of a substance 'X' is dissolved in 100 g of CCl_4 , it raises the boiling point by 0.60 K. The molar mass of the substance 'X' is _____ g mol^{-1} . (Nearest integer).
[Given K_b for CCl_4 is $5.0 \text{ K kg mol}^{-1}$]
16. 1.46 g of a biopolymer dissolved in a 100 mL water at 300 K exerted an osmotic pressure of $2.42 \times 10^{-3} \text{ bar}$.
The molar mass of the biopolymer is _____ $\times 10^4 \text{ g mol}^{-1}$.
(Round off to the Nearest Integer)
[Use : $R = 0.083 \text{ L bar mol}^{-1} \text{ K}^{-1}$]
17. When 400 mL of 0.2M H_2SO_4 solution is mixed with 600 mL of 0.1 M NaOH solution, the increase in temperature of the final solution is _____ $\times 10^{-2} \text{ K}$. (Round off to the nearest integer).
[Use : $\text{H}^+(\text{aq}) + \text{OH}^-(\text{aq}) \rightarrow \text{H}_2\text{O}$:
 $\Delta_r H = -57.1 \text{ kJ mol}^{-1}$
Specific heat of $\text{H}_2\text{O} = 4.18 \text{ J K}^{-1} \text{ g}^{-1}$
density of $\text{H}_2\text{O} = 1.0 \text{ g cm}^{-3}$
Assume no change in volume of solution on mixing.
18. Of the following four aqueous solutions, total number of those solutions whose freezing point is lower than that of 0.10 M $\text{C}_2\text{H}_5\text{OH}$ is _____ (Integer answer)
(i) 0.10 M $\text{Ba}_3(\text{PO}_4)_2$
(ii) 0.10 M Na_2SO_4
(iii) 0.10 M KCl
(iv) 0.10 M Li_3PO_4
19. 83 g of ethylene glycol dissolved in 625 g of water. The freezing point of the solution is _____ K. (Nearest integer)
[Use : Molal Freezing point depression constant of water = $1.86 \text{ K kg mol}^{-1}$
Freezing Point of water = 273 K
Atomic masses : C : 12.0 u, O : 16.0 u, H : 1.0 u]
20. 1 kg of 0.75 molal aqueous solution of sucrose can be cooled up to -4°C before freezing. The amount of ice (in g) that will be separated out is _____. (Nearest integer)
[Given : $K_f(\text{H}_2\text{O}) = 1.86 \text{ K kg mol}^{-1}$]
21. 40 g of glucose (Molar mass = 180) is mixed with 200 mL of water. The freezing point of solution is _____ K. (Nearest integer)
[Given : $K_f = 1.86 \text{ K kg mol}^{-1}$; Density of water = 1.00 g cm^{-3} ; Freezing point of water = 273.15 K]
22. Which one of the following 0.10 M aqueous solutions will exhibit the largest freezing point depression ?
(1) hydrazine (2) glucose
(3) glycine (4) KHSO_4
23. 1.22 g of an organic acid is separately dissolved in 100 g of benzene ($K_b = 2.6 \text{ K kg mol}^{-1}$) and 100 g of acetone ($K_b = 1.7 \text{ K kg mol}^{-1}$). The acid is known to dimerize in benzene but remain as a monomer in acetone. The boiling point of the solution in acetone increases by 0.17°C . The increase in boiling point of solution in benzene in $^{\circ}\text{C}$ is $x \times 10^{-2}$. The value of x is _____. (Nearest integer)
[Atomic mass : C = 12.0, H = 1.0, O = 16.0]

SOLUTION

1. Official Ans. by NTA (36)



$$i = 1 + (2 - 1)\alpha$$


$$i = 1 + \alpha$$

$$\Delta T_f = i k_f m$$

$$0.5 = (1 + \alpha)(1.86) \left(\frac{\left(\frac{9.45}{94.5} \right)}{\left(\frac{500}{1000} \right)} \right)$$

$$\frac{5}{3.72} = 1 + \alpha \Rightarrow \alpha = \frac{1.28}{3.72}$$

$$\alpha = \frac{32}{93}$$



$$K_a = \frac{(\text{C}\alpha)^2}{\text{C} - \text{C}\alpha} = \frac{\text{C}\alpha^2}{1 - \alpha} \quad \text{C} = \frac{0.1}{500/1000} = 0.2$$

$$K_a = \frac{0.2(32/93)^2}{(1 - 32/93)} = \frac{0.2 \times (32)^2}{93 \times 61}$$

$$= 0.036$$

$$K_a = 36 \times 10^{-3}$$

2. Official Ans. by NTA (1)

Sol. Pure Solvent : C_6H_6 (l)

$$\text{Given : } T_f^\circ = 5.5^\circ\text{C}$$

$$K_f = 5.12^\circ\text{C/m}$$



10g : Solute is non dissociative

200 g C_6H_6

$$\therefore \Delta T_f = k_f \times m$$

$$\Rightarrow (T_f^\circ - T_f') = 5.12 \times \frac{\left(\frac{10}{58} \right)}{\left(\frac{200}{1000} \right)} \text{ mol/kg}$$

$$\Rightarrow 5.5 - T_f' = \frac{5.12 \times 5 \times 10}{58}$$

$$\Rightarrow T_f' = 1.086^\circ\text{C} \approx 1^\circ\text{C}$$

3. Official Ans. by NTA (375)

Sol. $\Delta T_b = i K_b m$

$$= (1 + 4\alpha) \times 0.52 \times 1$$

$$= 3.4 \times 0.52 \times 1 = 1.768$$

$$T_b = 1.768 + 373.15 = 374.918 \text{ K}$$

$$= 375\text{K}$$

Hence answer is (375)

4. Official Ans. by NTA (3)

Sol. $\alpha = 0.75, n = 2$

$$i = 1 - \alpha + n\alpha = 1 - 0.75 + 2 \times 0.75 = 1.75$$


$$\Delta T_b = i k_b m$$

$$\text{or, } 2.5 = 1.75 \times 0.52 \times m$$

$$\text{or, } m = \frac{2.5}{1.75 \times 0.52} = 2.74$$

\therefore nearest integer answer will be 3

5. Official Ans. by NTA (24)



$$\frac{224}{0.0821 \times 298} \text{ (L.R.)} \quad \frac{10\text{mmol}}{(i = 3)} \quad \frac{5\text{mmol}}{(i = 3)}$$


$$= 9.2 \text{ m mol}$$

$$P^s = P^0 \cdot X_{\text{solvent}}$$

$$= 24 \times \frac{2}{(2 + 15 \times 10^{-3})}$$

$$= 23.82$$

$$\Delta P = 0.18 \text{ torr} = 18 \times 10^{-2} \text{ torr.}$$



$$\begin{array}{ccc} 9.2 & 10 & - \\ - & 0.8 & 9.2 \end{array}$$

$$\Delta P = P^0 \cdot X_{\text{solute}}$$

$$= 24 \times \frac{(1.6 + 18.4)}{2020}$$

$$= 0.2376 = 23.76 \times 10^{-2}$$

12. Official Ans. by NTA (100)

Sol. $\Delta T_b = T_b - T_b^0$

$$100.52 - 100$$

$$= 0.52^\circ\text{C}$$

$$i = \left(1 - \frac{\alpha}{2}\right)$$

$$\therefore \Delta T_b = i K_b \times m$$

$$0.52 = \left(1 - \frac{\alpha}{2}\right) \times 0.52 \times 2$$

$$\alpha = 1$$

So, percentage association = 100%.

13. Official Ans. by NTA (1)

Sol. $T_f - T_f' = i K_f \cdot m$

For minimum T_f'

'i' should be maximum.



14. Official Ans. by NTA (25)

Sol. From Henry's law

$$P_{\text{gas}} = K_H \cdot X_{\text{gas}}$$

$$0.835 = 1.67 \times 10^3 \times \frac{n(\text{CO}_2)}{0.9 \times 1000}$$

$$n(\text{CO}_2) = 0.025$$

$$\text{Millimoles of CO}_2 = 0.025 \times 1000 = 25$$

15. Official Ans. by NTA (250)

Sol. $\Delta T_b = K_b \times \text{molality}$

$$0.60 = 5 \times \left(\frac{3/M}{100/100}\right)$$

$$M = 250$$

16. Official Ans. by NTA (15)

Sol. $\pi = CRT$; π = osmotic pressure

$$C = \text{molarity}$$

$$T = \text{Temperature of solution}$$

let the molar mass be M gm / mol

$$2.42 \times 10^{-3} \text{ bar}$$

$$= \frac{\left(\frac{1.46 \text{ g}}{M \text{ gm / mol}}\right)}{0.1 \ell} \times \left(\frac{0.083 \ell - \text{bar}}{\text{mol} - \text{K}}\right) \times (300 \text{ K})$$

$$\Rightarrow M = 15.02 \times 10^4 \text{ g/mol}$$

17. Official Ans. by NTA (2)

ALLEN Ans. (82)

Sol. $n_{\text{H}^+} = \frac{400 \times 0.2}{1000} \times 2 = 0.16$

$$n_{\text{OH}^-} = \frac{600 \times 0.1}{1000} = 0.06 \text{ (L.R)}$$

Now, heat liberated from reaction

= heat gained by solutions

$$\text{or, } 0.06 \times 57.1 \times 10^3$$

$$= (1000 \times 1.0) \times 4.18 \times \Delta T$$

$$\therefore \Delta T = 0.8196 \text{ K}$$

$$= 81.96 \times 10^{-2} \text{ K} \approx 82 \times 10^{-2} \text{ K}$$

18. Official Ans. by NTA (4)


Sol. As 0.1 M $\text{C}_2\text{H}_5\text{OH}$ is non-dissociative and rest all salt given are electrolyte so in each case effective molarity > 0.1 so each will have lower freezing point.

19. Official Ans. by NTA (269)

Sol. $k_f = 1.86 \text{ k. kg/mol}$

$$T_f^0 = 273 \text{ K}$$

solvent : H_2O (625 g)



solute

$$\Rightarrow \Delta T_f = k_f \times m$$

$$\Rightarrow (T_f^0 - T_f^1) = 1.86 \times \frac{83 / 62}{624 / 1000}$$

$$\Rightarrow 273 - T_f^1 = \frac{1.86 \times 83 \times 1000}{62 \times 625} = \frac{154380}{38750}$$

$$\Rightarrow 273 - T_f^1 = 4$$

$$\Rightarrow \boxed{T_f^1 = 259 \text{ K}}$$

20. Official Ans. by NTA (518)

Sol. Let mass of water initially present = x gm

$$\Rightarrow \text{Mass of sucrose} = (1000 - x) \text{ gm}$$

$$\Rightarrow \text{moles of sucrose} = \left(\frac{1000 - x}{342} \right)$$

$$\Rightarrow 0.75 = \frac{\left(\frac{1000 - x}{342} \right)}{\left(\frac{x}{1000} \right)} \Rightarrow \frac{x}{1000} = \frac{1000 - x}{342 \times 0.75}$$

$$\Rightarrow 256.5x = 10^6 - 1000x$$

$$\Rightarrow x = 795.86 \text{ gm}$$

$$\Rightarrow \text{moles of sucrose} = 0.5969$$

New mass of H_2O = a kg

$$\Rightarrow 4 = \frac{0.5969}{a} \times 1.86 \Rightarrow a = 0.2775 \text{ kg}$$

$$\Rightarrow \text{ice separated} = (795.86 - 277.5) = 518.3 \text{ gm}$$

21. Official Ans. by NTA (271)

Sol. molality = $\frac{\left(\frac{40}{180} \right) \text{ mol}}{0.2 \text{ Kg}} = \left(\frac{10}{9} \right) \text{ molal}$

$$\Rightarrow \Delta T_f = T_f - T_f' = 1.86 \times \frac{10}{9}$$

$$\Rightarrow T_f' = 273.15 - 1.86 \times \frac{10}{9}$$

$$= 271.08 \text{ K}$$

$$\approx 271 \text{ K (nearest-integer)}$$

22. Official Ans. by NTA (4)

Sol. \therefore Van't Hoff factor is highest for KHSO_4

\therefore colligative property (ΔT_f) will be highest for KHSO_4

23. Official Ans. by NTA (13)

Sol. With benzene as solvent

$$\Delta T_b = i K_b m$$

$$\Delta T_b = \frac{1}{2} \times 2.6 \times \frac{1.22 / M_w}{100 / 1000} \quad \dots(1)$$

With Acetone as solvent

$$\Delta T_b = i K_b m$$

$$0.17 = 1 \times 1.7 \times \frac{1.22 / M_w}{100 / 1000} \quad \dots(2)$$

(1) / (2)

$$\frac{\Delta T_b}{0.17} = \frac{\frac{1}{2} \times 2.6 + \frac{1.22 / M_w}{100 / 1000}}{1 \times 1.7 \times \frac{1.22 / M_w}{100 / 1000}}$$

$$\Delta T_b = \frac{0.26}{2}$$

$$\Delta T_b = 13 \times 10^{-2}$$

$$\Rightarrow x = 13$$